

# Swan Canning Estuary condition assessment based on fish communities - 2022

James Tweedley, Kurt Krispyn & Alan Cottingham

*Centre for Sustainable Aquatic Ecosystems, Harry Butler Institute, Murdoch University  
College of Science, Health, Engineering & Education, Murdoch University*

## Final Report

Prepared for the Department of Biodiversity, Conservation and Attractions



Department of **Biodiversity,  
Conservation and Attractions**



## **Disclaimer**

The authors have prepared this report in accordance with the scope of work and for the purpose required by the Department of Biodiversity, Conservation and Attractions. The methodology adopted and sources of information used by the author are outlined in this report. The authors have made no independent verification of this information beyond the agreed scope of works and assumes no responsibility for any inaccuracies or omissions. The authors disclaim any responsibility for changes that may have occurred after this time. This report should be read in full. No responsibility is accepted for use of any part of this report in any other context or for any other purpose or by third parties. This report does not purport to give legal advice. Legal advice can only be given by qualified legal practitioners.

## **Acknowledgements**

Gratitude is expressed to Jake Watsham and Sarah Muller from the Department of Biodiversity, Conservation and Attractions who assisted with the sampling of the fish community. Dr Steeg Hoeksema is thanked for helping to coordinate the sampling program, Dr Jeff Cosgrove for supplying information on phytoplankton composition and densities and Dr Kerry Trayler for reviewing the draft final report.

## **Recommended citation:**

Tweedley, J.R., Krispyn, K.N. & Cottingham, A. (2022). Swan Canning Estuary condition assessment based on fish communities - 2022. Final report to the Department of Biodiversity, Conservation and Attractions. Murdoch University, Western Australia, 59 pp.

*Cover image acknowledgements: Kurt Krispyn, Matt Hourston, Dave Morgan, Mark Allen, Gavin Sarre (Murdoch University)*

## Executive summary

This report, commissioned by the Department of Biodiversity, Conservation and Attractions (DBCA), describes the monitoring and evaluation of fish communities in the Swan Canning Estuary during 2022 and applies the Fish Community Index (FCI) that was developed as a measure of the ecological condition of the Swan Canning Estuary. This index, separate versions of which were developed for both the shallow (< 1.5 m), nearshore waters of the estuary and also for its deeper (> 1.5 m), offshore waters, integrates information on various biological variables (metrics). Each of these metrics quantifies an aspect of the structure and/or function of the fish community, and together they respond to a range of stressors affecting the ecosystem.

Fish communities were sampled using different types of net at six nearshore and six offshore sites in each of four management zones of the estuary (LSCE, Lower Swan Canning Estuary; CE, Canning Estuary; MSE, Middle Swan Estuary; USE, Upper Swan Estuary) during summer and autumn of 2022. As many fish as possible were returned to the water alive after they had been identified and counted. The resulting data on the abundances of each fish species from each sample were used to calculate a Fish Community Index score (0–100). These index scores were then compared to established scoring thresholds to determine ecological condition grades (A–E) for each zone and for the estuary as a whole, based on the composition of the fish community.

### *Nearshore Fish Communities*

The nearshore waters of the estuary as a whole were in good (B) and good/fair condition (B/C) during summer and autumn 2022, respectively, which is consistent with the overall trend in condition since 2011. The average nearshore FCI scores for each zone of the estuary varied during summer, being best, i.e. good, in the LSCE, MSE and USE and lowest in the CE resulting in a fair (C) score. By autumn, scores in the CE increased to good, while those in the LSCE and MSE declined slightly from good to good/fair in the LSCE and good to fair in the MSE. While the mean score for the USE dropped the grade remained the same, i.e. good. There were no notable algal blooms in the MSE or USE and although some stratification was present, oxygen concentrations did not decline sufficiently to cause hypoxia.

Small-bodied, schooling species of hardyheads (Atherinidae) and gobies (Gobiidae) once again dominated catches from the nearshore waters of the estuary in 2021, representing 77% of all fish recorded and constituting the six of the seven most abundant nearshore species. Wallace's Hardyhead was the most abundant species overall and also in the CE and USE, reflecting the preference of this species for the fresh to brackish conditions that were present in these zones during the 2022 monitoring period. Juvenile Western Striped Grunter made an unusually large contribution to the fish fauna, driven by good recruitment from the ocean where they spawn. Other abundant species of small, schooling fish included the Spotted Hardyhead, Common Hardyhead and Silverfish, each of which prefer more saline waters than Wallace's Hardyhead. The Perth Herring was abundant in the MSE and USE, as were the Bluespot Goby, Black Bream and the Australian Anchovy in the USE.

The total number of fish caught in nearshore waters in 2022 was 1.4 to 2.6 times more than that recorded previously (2012-2021). This marked increase in the density of fish is likely the result of flows from the Swan River in 2021 being the largest for 25 years. Strong flows are known to increase productivity and fish populations. Stable salinities and oxic conditions during the 2022 monitoring

period, combined with good recruitment, resulted in the largest number of species (36) being recorded.

### *Offshore fish communities*

The offshore waters of the Swan Canning Estuary were in good (B) condition in summer and fair/good (C/B) during autumn 2022. The overall score of good was an increase on the fair (C) grade it received in 2021 and represents only the fourth time in 14 years good condition was achieved. Scores in the LSCE, MSE and USE were either very good (A) or good, except for fair/good in the MSE in autumn, likely driven by relatively saline and oxic conditions and the absence of toxic algal blooms. However, once again the offshore waters of the CE exhibited by far the lowest scores of any zone. As the Canning River is more regulated than the Swan, for example by the presence of the Canning Dam, this axis of the estuary did not receive the substantial increase in flow that the USE, MSE and to some extent the LSCE did. Waters in the CE were stratified, hypoxic, and at times, contained the toxic dinoflagellate *Karlodinium* spp..

As in most previous years of monitoring, Perth Herring was among the dominant species in offshore waters from all four zones comprising 29–80% of the total catches. Other abundant species included the Southern Eagle Ray and Tailor in the LSCE (29 and 10%, respectively, of the catch) and the Yellowtail Grunter in the MSE (24%) and USE (38%) and Black Bream in the USE (5%). The number of species and individuals recorded from the offshore waters in 2022 were amongst the greatest in any monitoring year. Catches of several species were relatively high in 2022, including the Southern Eagle Ray, Mulloway and Australian Giant Herring.

### *Overall*

Across the entire estuary, the ecological condition of both nearshore and offshore waters in 2022 was assessed as good (B) based on their fish communities. Combined, the nearshore and offshore index scores for 2022 are the highest ever recorded since annual monitoring began in 2012. The good scores for zones along the Swan axis were likely influenced by the strong freshwater flows that occurred in the winter of 2021. Such flows, which were the greatest for 25 years, increased productivity in both the estuary and also nearby coastal waters and facilitated the recruitment of fish species. Moreover, the absence of atypical summer rainfall events and low phytoplankton densities in most regions facilitated the maintenance of relatively saline and oxic conditions in most zones except the CE where some stratification-induced hypoxia occurred and the toxic dinoflagellate *Karlodinium* spp. was detected.

# Swan Canning Estuary condition assessment based on fish communities - 2022

## Contents

Acknowledgements.....	2
Executive summary .....	3
Contents.....	5
1. Background .....	6
2. Rationale .....	6
3. Study objectives .....	7
4. Methods .....	8
5. Results and discussion .....	11
5.1 Context: water quality and environmental conditions during the 2022 monitoring period .....	11
5.2 The fish community of the Swan Canning Estuary during 2022 .....	13
Nearshore waters.....	13
Offshore waters .....	17
5.3 Ecological condition in 2022 .....	19
Nearshore waters.....	19
Offshore waters .....	21
Longer term trends in ecological condition .....	24
Summary .....	26
6. References .....	27
7. Appendices.....	31

## 1. Background

The Department of Biodiversity, Conservation and Attractions (DBCA) works with other government organizations, local government authorities, community groups and research institutions to reduce nutrient and organic loading to the Swan Canning Estuary and river system. This is a priority issue for the waterway that has impacts on water quality, ecological health and community benefit. Environmental monitoring for the waterway includes water quality reporting in the estuary and catchment and reporting on ecological health. Reporting on changes in fish communities provides insight into the biotic integrity of the system and complements water quality reporting.

The Fish Community Index (FCI) was developed by Murdoch University, in collaboration with the Western Australian government between 2007 and 2012 (Valesini et al., 2011; Hallett and Valesini, 2012; Hallett et al., 2012), and provides an assessment the condition of the Swan Canning Estuary based on fish communities. The FCI has been subjected to extensive testing and validation over a period of many years (e.g. Hallett and Valesini, 2012; Hallett, 2014), and has been shown to be a sensitive and robust tool for quantifying ecological health responses to local-scale environmental perturbations and the subsequent recovery of the system following their removal (Hallett, 2012; Hallett et al., 2012; 2016). The development and rationale of the FCI, along with its implementation and outcomes to date, are summarized in Hallett et al. (2019).

## 2. Rationale

Separate versions of the FCI were developed for the shallow, nearshore waters (< 1.5 m deep) of the estuary and also for its deeper, offshore waters (> 1.5 m deep), as the composition of the fish communities living in these different environments tends to differ, as do the methods used to sample them (Chuwen, 2009; Hoeksema et al., 2009; Potter et al., 2016). These indices integrate information on various biological variables ('metrics'; Table 1), each of which quantifies an aspect of the structure and/or function of estuarine fish communities. Together, the metrics respond to a wide array of stressors affecting the ecosystem. The FCI therefore provides a means to assess an important component of the ecology of the system and how it responds to, and thus reflects, changes in estuarine condition (Hallett et al., 2019; Tweedley et al., 2021).

The responses of estuarine fish communities to increasing ecosystem stress and degradation (i.e. declining ecosystem health or condition) may be summarised in a conceptual model (Fig. 1). In response to increasing degradation of estuarine ecosystems, fish species with specific habitat, feeding or other environmental requirements will tend to become less abundant and diverse, whilst a few species with more general requirements become more abundant. This leads ultimately to an overall reduction in the number and diversity of fish species (Gibson et al., 2000; Whitfield and Elliott, 2002; Villéger et al., 2010; Fonseca et al., 2013; Tweedley et al., 2017). So, in a degraded estuary with poor water, sediment and habitat quality, the abundance and diversity of specialist feeders (e.g. Garfish and Tailor), bottom-living ('benthic-associated') species (e.g. Cobbler and Flathead) and estuarine spawning species (e.g. Black Bream, Perth Herring and Yellowtail Grunter) will tend to decrease, as will the overall number and diversity of species. In contrast, generalist feeders (e.g. Banded Toadfish or Blowfish) and detritivores (e.g. Sea Mullet), which eat particles of decomposing organic material, will become more abundant and dominant (Krispyn et al., 2021; right side of Fig. 1). The reverse will be observed in a relatively unspoiled system that is subjected to fewer human stressors (see left side of Fig. 1; noting that this conceptual diagram represents either end of a continuum of ecological condition from very poor to very good).

Each of the metrics that make up the FCI are scored from 0–10 according to the numbers and proportions of the various fish species present in samples collected from the estuary using either seine or gill nets. These metric scores are summed to generate an FCI score for the sample, which ranges from 0–100. Grades (A–E) describing the condition of the estuary, and/or of particular zones, are then awarded based on the FCI scores (see Section 4 for more details).

**Table 1.** Summary of the metrics comprising the nearshore and offshore Fish Community Indices developed for the Swan Canning Estuary (Hallett et al., 2012).

<i>Metric</i>	<b>Predicted response to degradation</b>	<i>Nearshore Index</i>	<i>Offshore Index</i>
Number of species (No. species)	Decrease	√	√
Shannon-Wiener diversity (Sh-div) <sup>a</sup>	Decrease		√
Proportion of trophic specialists (Prop. trop. spec.) <sup>b</sup>	Decrease	√	
Number of trophic specialist species (No. trop. spec.) <sup>b</sup>	Decrease	√	√
Number of trophic generalist species (No. trop. gen.) <sup>c</sup>	Increase	√	√
Proportion of detritivores (Prop. detr.) <sup>d</sup>	Increase	√	√
Proportion of benthic-associated individuals (Prop. benthic) <sup>e</sup>	Decrease	√	√
Number of benthic-associated species (No. benthic) <sup>e</sup>	Decrease	√	
Proportion of estuarine-spawning individuals (Prop. est. spawn)	Decrease	√	√
Number of estuarine-spawning species (No. est. spawn)	Decrease	√	
Proportion of <i>Pseudogobius olorum</i> (Prop. <i>P. olorum</i> ) <sup>f</sup>	Increase	√	
Total number of <i>Pseudogobius olorum</i> (Tot no. <i>P. olorum</i> ) <sup>f</sup>	Increase	√	

<sup>a</sup> A measure of biodiversity

<sup>b</sup> Species with specialist feeding requirements (e.g. those that only eat small invertebrates)

<sup>c</sup> Species that are omnivorous or opportunistic feeders

<sup>d</sup> Species that eat detritus (decomposing organic material)


<sup>e</sup> Species that live on, or are closely associated with, the sea/river bed

<sup>f</sup> The Bluespot or Swan River Goby, a tolerant, omnivorous species that often inhabits silty habitats (Gill and Potter, 1993)

### 3. Study objectives

This report describes the monitoring and evaluation of fish communities in the Swan Canning Estuary during 2022 for the purposes of applying the FCI as a measure of ecological condition. The objectives of this study were to:

1. Undertake monitoring of fish communities in mid-summer and mid-autumn periods, following an established approach as detailed in Hallett and Valesini (2012), including six nearshore and six offshore sampling sites in each estuarine management zone.
2. Analyse the information collected so that the FCI is calculated for nearshore and offshore waters in each management zone and for the estuary overall. The information shall be presented as quantitative FCI scores (0–100), qualitative condition grades (A–E) and descriptions of the fish communities. Radar plots shall also be used to demonstrate the patterns of metric scores for each zone.
3. Provide a report that summarizes the approach and results and that could feed into the broader estuarine reporting framework of the Department of Biodiversity, Conservation and Attractions.



**Figure 1.** Conceptual diagram illustrating the predicted responses of the estuarine fish community to situations of very good (A) and very poor (E) ecological condition. Images courtesy of the Integration and Application Network [ian.umces.edu/symbols/].

#### 4. Methods

Fish communities were sampled at six nearshore and six offshore sites in each of the four management zones of the Swan Canning Estuary (LSCE, Lower Swan Canning Estuary; CE, Canning Estuary; MSE, Middle Swan Estuary; USE, Upper Swan Estuary; Fig. 2) during both summer (31 January – 17 February) and autumn (19 April – 4 May) of 2022. All sampling was conducted under permits approved by Murdoch University’s Animal Ethics Committee (permit number RW3286/20), the Department of Primary Industries and Regional Development, Fisheries Division (exemption number 3585) and Department of Biodiversity, Conservation and Attractions (permit number FO25000254-3).

Nearshore waters were sampled using a 21.5 m seine net that was walked out from the beach to a maximum depth of ~ 1.5 m and deployed parallel to the shore, and then rapidly dragged towards


and onto the shore (Fig. 3). Offshore waters were sampled using 160 m-long, sunken, multimesh gill nets, each consisting of eight 20 m-long panels with stretched mesh sizes of 35, 51, 63, 76, 89, 102, 115 and 127 mm (Fig. 3). These were deployed (i.e. laid parallel to the bank at a depth of 2–8 m, depending on the depth of water at each site) from a boat immediately before sunset and retrieved after three hours.

Once a sample had been collected, any fish that could be identified immediately to species (e.g. larger species that are caught in relatively lower numbers) were identified, counted and returned to the water alive. All other fish caught in the nets were placed into zip-lock polythene bags, euthanised in ice slurry and preserved on ice for subsequent identification and counting, except in cases where large catches (e.g. thousands) of small fish were obtained. In such instances, an appropriate sub-sample (e.g. one-half to one-eighth of the catch, depending on the total size of the catch) was retained for identification and estimation of the numbers of each species, and the remaining fish were returned alive to the water to minimise the impact on fish populations. All retained fish were then frozen until their identification in the laboratory by experienced fish biologists, using available keys and identification guides where required. See appendices (i and ii) for full details of the sampling locations and methods employed.

The abundances of each fish species in each sample were used to derive values for each of the relevant metrics comprising the nearshore and offshore indices (Hallett and Valesini, 2012; Hallett et al., 2012) using bespoke code developed for the R software package. Metric scores were then calculated from these metric values, and the metric scores in turn combined to form the FCI scores. The method for calculating these scores is detailed in Hallett and Valesini (2012), but can be summarised simply as follows:

1. Allocate each fish species in a particular sample to its appropriate Habitat guild, Estuarine Use guild and Feeding Mode guild (Appendix iii), then calculate the values for each fish metric from the abundances of fishes in the sample.
2. Convert metric values to metric scores (0–10) via comparison with the relevant (zone- and season-specific) reference condition values for each metric.
3. Combine scores for the component metrics into a scaled FCI score (0–100) for each sample.
4. Compare the FCI score to the thresholds used to determine the condition grade for each sample (Table 2; Hallett, 2014), noting that intermediate grades e.g. B/C (good/fair) or C/B (fair/good) are awarded if the index score lies within one point either side of a grade threshold.


The FCI scores and condition grades for nearshore and offshore samples collected during summer and autumn 2022 were then examined to assess the condition of the Swan Canning Estuary during this period and were compared to previous years through a qualitative examination of the patterns and trends in scores.



**Figure 2.** Locations of nearshore (light blue circles) and offshore (dark blue circles) sampling sites for the Fish Community Index of estuarine condition.

**Table 2.** Fish Community Index (FCI) scores comprising each of the five condition grades for both nearshore and offshore waters of the Swan Canning Estuary. Intermediate grades, e.g. B/C (good/fair) or C/B (fair/good) are awarded if the index score lies within one point either side of a grade threshold.

Condition grade	Nearshore FCI scores	Offshore FCI scores
A (very good)	> 74.5	> 70.7
B (good)	64.6 - 74.5	58.4 - 70.7
C (fair)	57.1 - 64.6	50.6 - 58.4
D (poor)	45.5 - 57.1	36.8 - 50.6
E (very poor)	< 45.5	< 36.8



**Figure 3.** Photographs of the beach seine netting (upper row) used to sample the fish community in shallow, nearshore waters and the multimesh gill netting (lower row) used to sample fish communities in deeper, offshore waters of the Swan Canning Estuary. Images courtesy of Kurt Krispyn, Murdoch University.

## 5. Results and discussion

### 5.1 Context: water quality and environmental conditions during the 2022 monitoring period

Total annual flow at Walyunga on the Swan River in 2021 was 605 GL, the highest recorded since 1996 some 25 years earlier and sixth highest since records began in 1971 and 303% increase over the average between 1996 and 2020 (Appendix iv). The timing of the flow corresponded with the traditional monthly pattern, where the majority of flow occurs between May and September (Hodgkin and Hesp, 1998; Hallett et al., 2018). In 2022, 92% of the total annual flow occurred between these months, with particularly high values recorded in July and August (235 and 248 GL, respectively). The volume of flow in each of these two months was the greatest since February 2017 (Appendix iv), following one of the wettest days recorded in Perth and that resulted in widespread hypoxia and one of the lowest offshore FCI scores (Hallett, 2017).

Total annual flow at Seaforth in the Canning River in 2021 was 10.4 GL, with 39% occurring in July, which is over twice the average for this month (Appendix v). Although the volume of flow in the Canning River in 2021 was 24% greater than the median (~8.4 GL) recorded since 1996, this increase was far less than that recorded at Walyunga. This marked difference in the increases of flow between the Swan and Canning rivers in 2021 compared to the longer-term is likely influenced by the fact that the Canning is heavily regulated by two major water supply dams, (Canning Reservoir and Wungong

Reservoir), two smaller dams (Churchman Brook Reservoir and Victoria Reservoir) and a number of smaller weirs (Radin et al., 2007; Norton et al., 2010).

The environmental conditions present in the system during the monitoring period are shown as vertical contour plots of interpolated salinities, dissolved oxygen (DO) concentrations, chlorophyll levels and water temperatures measured at regular water quality monitoring sites along the length of both the Swan and Canning axes of the estuary on a weekly basis (Appendix vi). The water column of the USE was brackish (salinity 6 - 18) in early January 2021, becoming more saline into mid-February (minimum of 12) as the salt wedge moved upstream. Salinities in the LSCE and MSE were around that of full-strength sea water (~35) throughout summer ranging from 30 to 40. Areas of low dissolved oxygen (2 - 4 mg/L) and smaller pockets of hypoxia (< 2 mg/L) occurred in parts of the MSE and USE during some weeks in January and February. Both the Caversham and Guildford Oxygenation plants were in operation in each week of January and February. Water temperature increased in an upstream direction from ~22 °C in the LSCE to 27 - 29 °C in the USE. In terms of phytoplankton, densities were low during the summer sampling period in the LCSE, with moderate levels of the dinoflagellates *Heterocapsa* spp. (5,300 cells/mL), *Karlodinium* spp. (5,400 cells/mL) and *Gymnodinium-Karenia* Complex (1,200 cell/mL) with values of the first two taxa being recorded in greater densities at Ron Courtney Island at the lower end of the USE (i.e. 7,760 and 5,240 cells/mL, respectively; DBCA, unpublished data). Densities of the dinoflagellates *Heterocapsa minima* and *Heterocapsa lanceolata* at Meadow Street in the USE exceeded management trigger values, but this event occurred towards the very end of the sampling program in summer and so would have been unlikely to have influenced the results.

The water column of the upper part of the CE (Riverton to Castledare) was stratified by freshwater flows overlying denser, saltier water in January and February with the degree of stratification decreasing over time as more of this zone became saline (Appendix vi). Pronounced stratification was detected on 25 January (immediately prior to the initiation of sampling) and again two weeks later on 8 February, resulting in the presence of hypoxic conditions (< 2 mg/L) in these weeks. Phytoplankton densities were moderate through January and early February in the CE, with the highest densities of > 30,000 cells/mL recorded on 25 January at Shelley Riverton and Castledare in the upper CE. In these samples, moderate levels of densities of *Karlodinium* spp. (8,500 cells/mL) and *Heterosigma akashiwo* (17,380 cells/mL) were recorded (DBCA, unpublished data).

During autumn sampling (19 April – 4 May), marine conditions, i.e. salinities of ~35, were found throughout the LSCE and salinities decreased in an upstream direction to a minimum of ~20 in the upper parts of the USE (Appendix vi). The waters of the MSE and USE were stratified, caused by the salt wedge at the bottom of the water column moving upstream during these months, however, this did not lead to the development of hypoxia. The absence of low dissolved oxygen concentrations could also be due to the continued operation of the Guildford and Caversham oxygenation plants during the sampling period. Total phytoplankton densities were low to moderate with peak densities across the estuary recorded at Ron Courtney Island (16,800 cells/mL) and West Midland Pool (17,725 cells/mL) in the USE on 19 and 26 April (DBCA, unpublished data). Densities declined substantially in this zone in early May, with the highest value in the estuary of 12,700 cells/mL being recorded at Maylands in the MSE. No blooms of harmful species were detected over the sampling period.

Environmental conditions in the CE during autumn were affected by the pronounced stratification present in the upper parts of the CE, which resulted in hypoxia. This hypoxia lasted until ~10 May and thus after sampling was completed. Total phytoplankton densities were low in the lower

reaches of the CE (Canning Bridge and Salter Point) but increased to moderate levels at Riverton and Castledare. There was no evidence of harmful blooms (DBCA, unpublished data).

## **5.2 The fish community of the Swan Canning Estuary during 2022**

### *Nearshore waters*

An estimated total of 42,935 fish, belonging to 36 species, were caught in seine net samples collected from the nearshore waters during the summer and autumn of 2022. The total number of fish caught in that year (i.e. 42,935) was 1.4 to 2.6 times more than recorded in any years previously (range = 16,905 - 30,825). This marked increase in the density of fish is likely the result of the increased riverine flows in 2022, which are known to increase productivity and fish populations (Gillanders and Kingsford, 2002; Broadley et al., 2022). The 36 species recorded in 2022 was slightly greater than the 35 and 32 in 2021 and 2020 and above the annual average of 31.8 (range = 25 – 35). A total of 64 fish species have been collected in seine nets as part of this annual monitoring since 2012. One new addition to the nearshore fish fauna was recorded in 2022, i.e. the Tailor (*Pomatomus saltatrix*), albeit larger-bodied individuals of this species are common in offshore waters. The greatest number of species recorded in the nearshore waters was in the LSCE and CE (both 25), followed by the MSE (24) and least in the USE (20; Table 3). This pattern of decline in the number of species along the longitudinal (downstream – upstream) axis has been recorded in the nearshore waters of Swan Canning previously and in similar estuaries in south-western Australia (Veale et al., 2014; Valesini et al., 2017). The total number of species in each zone except the LSCE, was significantly greater than those between 2012 and 2021. This can be explained by the presence of a wider range of marine-spawning species occurring further upstream than usual, such as Yellowtail Flathead (*Platycephalus westraliae*), Western Trumpeter Whiting (*Sillago berrus*) and Whitebait (*Hyperlophus vittatus*).

Hardyheads (family = Atherinidae; five species) and gobies (family = Gobiidae; seven species) once again dominated catches from the nearshore waters of the estuary in 2022, representing 77% of all fish caught and containing six of the seven most abundant nearshore species. In particular, Wallace's Hardyhead (*Leptatherina wallacei*) was again the most abundant species overall (Table 3) but with densities in 2022 being three times higher than the average, likely due to good recruitment following spawning at the end of 2021. Among zones of the estuary, this species ranked first in terms of density in the CE and USE, comprising 53 and 39 % of all fish, respectively, reflecting the preference of this species for upstream areas where salinities are less than in other parts of the estuary (Prince and Potter, 1983; Potter et al., 2015b). Another atherinid species, the Spotted Hardyhead (*Craterocephalus mugiloides*), which prefers slightly more saline waters than Wallace's Hardyhead, was amongst the most abundant species in the LSCE and amongst the most abundant in the MSE and CE. Together with *C. mugiloides* three other atherinids, Common Hardyhead (*Atherinomorus vaigiensis*) and Silverfish (*Leptatherina presbyteroides*) all of which prefer more saline waters dominated the fish found in the LSCE (Valesini et al., 2009; 2017). One species that made a large contribution to the fish fauna of the LSCE was the Western Striped Grunter (*Helotes octolineatus*) at 26%, furthermore, the abundance of this commonly-recorded species was eight times greater in 2022 than the average between 2012 and 2021. Individuals of this species in 2022 were small and likely new recruits spawned in the marine waters in spring (Veale et al., 2015). Other abundant species included the Bluespot goby (*Pseudogobius olorum*) in the USE and the Redspot Goby (*Favonigobius punctatus*) in the MSE where they typically occurs (Hogan-West et al., 2019), the Yelloweye Mullet (*Aldrichetta forsteri*) in the MSE, the Perth Herring (*Nematalosa vlaminghi*) in the MSE and USE and (Australian Anchovy (*Engraulis australis*) and Black Bream (*Acanthopagrus butcheri*) in the USE (Table 3).

Two non-native fish species were recorded namely the Eastern Gambusia (*Gambusia holbrooki*), which is known to act antagonistically to native species (Beatty et al., 2022), in the CE and USE as was the Pearl Cichlid (*Geophagus brasiliensis*) in the CE (Table 3). These species occur regularly in this annual monitoring program, being found in eleven and nine of the last eleven years, respectively. Numbers of the Eastern Gambusia caught in all samples collected in 2022 were the third highest recorded (i.e. 1,092; range = 37 – 1,633) and those of the Pearl Cichlid were the second greatest (i.e. 51; range = 0 – 60). The two recently recorded species of goby in the Swan Canning Estuary, i.e. the Largemouth Goby (*Redigobius macrostoma*) and Dusky Frillgoby (*Bathygobius fuscus*) were once again recorded in 2022. Despite the focused sampling regime employed in this study (i.e. only six samples per management zone), both species have been recorded in two of the last three years, which may indicate populations have become established. Future sampling will help confirm this hypothesis.

**Table 3.** Compositions of the fish communities (D = Average density fish/100 m<sup>2</sup> and %C = percentage composition) observed across the six nearshore sites sampled in each zone of the Swan Canning Estuary during summer and autumn of 2022. Data for the three most abundant species in the catches from each zone are emboldened for emphasis. Species ordered by total abundance throughout the estuary. LSCE = Lower Swan Canning Estuary, CE = Canning Estuary, MSE = Middle Swan Estuary, USE = Upper Swan Estuary. \* denotes non-native species.

Species	Common name	LSCE (n = 12)		CE (n = 12)		MSE (n = 12)		USE (n = 12)	
		D	%C	D	%C	D	%C	D	%C
<i>Leptatherina wallacei</i>	Wallace's Hardyhead	0.22	0.05	<b>1,209.27</b>	<b>53.23</b>	12.07	8.18	<b>75.43</b>	<b>38.65</b>
<i>Craterocephalus mugiloides</i>	Mugil's Hardyhead	<b>190.09</b>	<b>40.46</b>	<b>461.06</b>	<b>20.29</b>	<b>16.59</b>	<b>11.25</b>	0.93	0.48
<i>Helotes octolineatus</i>	Western Striped Grunter	<b>123.99</b>	<b>26.40</b>	<b>198.35</b>	<b>8.73</b>	<b>15.52</b>	<b>10.52</b>	2.37	1.21
<i>Atherinosoma elongatum</i>	Elongate Hardyhead	20.19	4.30	113.07	4.98	0.72	0.49		
<i>Favonigobius punctatus</i>	Red-spot Goby	15.23	3.24	62.14	2.74	11.85	8.03	10.27	5.26
<i>Atherinomorus vaigiensis</i>	Common Hardyhead	<b>51.01</b>	<b>10.86</b>	2.44	0.11	3.45	2.34	0.14	0.07
<i>Pseudogobius olorum</i>	Blue-spot / Swan River Goby			72.99	3.21	2.95	2.00	12.00	6.15
<i>Gambusia holbrooki</i> *	Mosquito Fish *			72.92	3.21			5.53	2.83
<i>Sillago burrus</i>	Western Trumpeter Whiting	23.06	4.91	13.00	0.57	9.77	6.62	0.65	0.33
<i>Nematalosa vlaminghi</i>	Perth Herring			5.24	0.23	10.63	7.21	<b>31.75</b>	<b>16.27</b>
<i>Leptatherina presbyteroides</i>	Presbyter's Hardyhead	16.02	3.41	1.08	0.05	0.36	0.24		
<i>Engraulis australis</i>	Southern Anchovy					0.72	0.49	<b>28.95</b>	<b>14.83</b>
<i>Acanthopagrus butcheri</i>	Southern Black Bream	0.14	0.03	8.55	0.38	9.41	6.38	11.28	5.78
<i>Amniataba caudavittata</i>	Yellow-tail Trumpeter	1.44	0.31	14.66	0.65	5.60	3.80	4.31	2.21
<i>Hyperlophus vittatus</i>	Whitebait / Sandy Sprat	9.70	2.06			3.88	2.63	1.87	0.96
<i>Aldrichetta forsteri</i>	Yellow-eye Mullet	0.36	0.08	4.31	0.19	<b>19.40</b>	<b>13.15</b>	0.14	0.07
<i>Favonigobius lateralis</i>	Long-finned Goby	8.19	1.74	4.60	0.20				
<i>Ostorhinchus rueppellii</i>	Gobbleguts	4.74	1.01			9.34	6.33		
<i>Gerres subfasciatus</i>	Roach	1.08	0.23	2.01	0.09	7.61	5.16	3.74	1.91
<i>Torquigener pleurogramma</i>	Blowfish / Banded Toadfish	0.65	0.14	7.83	0.34	1.58	1.07		
<i>Mugil cephalus</i>	Sea Mullet	0.43	0.09	4.17	0.18	3.23	2.19	1.29	0.66

Table 3. continued.

Species	Common name	LSCE (n = 12)		CE (n = 12)		MSE (n = 12)		USE (n = 12)	
		D	%C	D	%C	D	%C	D	%C
<i>Afurcagobius suppositus</i>	Southwestern Goby			3.45	0.15	0.65	0.44	3.81	1.95
<i>Geophagus brasiliensis</i> *	Pearl Cichlid *			3.16	0.14			0.50	0.26
<i>Arenigobius bifrenatus</i>	Bridled Goby			2.95	0.13	0.50	0.34	0.07	0.04
<i>Sillago schomburgkii</i>	Yellow-finned Whiting	0.29	0.06	2.30	0.10	0.50	0.34		
<i>Rhabdosargus sarba</i>	Tarwhine	1.65	0.35						
<i>Redigobius macrostoma</i>	Largemouth Goby			1.58	0.07				
<i>Pomatomus saltatrix</i>	Tailor					1.15	0.78		
<i>Haletta semifasciata</i>	Blue Weed Whiting	0.57	0.12						
<i>Gymnapistes marmoratus</i>	Devilfish	0.14	0.03	0.72	0.03				
<i>Bathygobius fuscus</i>	Dusky Frillgoby	0.29	0.06						
<i>Contusus brevicaudus</i>	Prickly Toadfish	0.14	0.03						
<i>Platycephalus westraliae</i>	Yellowtail Flathead					0.07	0.05	0.14	0.07
<i>Siphonognathus radiatus</i>	Long-rayed Weed Whiting	0.07	0.02						
<i>Scobinichthys granulatus</i>	Rough Leatherjacket	0.07	0.02						
<i>Pseudocaranx wrightii</i>	Skipjack Trevally			0.07	0.00				
<b>Total number of species</b>		<b>25</b>		<b>25</b>		<b>24</b>		<b>20</b>	
<b>Average total fish density (fish 100m<sup>-2</sup>)</b>		<b>470</b>		<b>2272</b>		<b>148</b>		<b>195</b>	
<b>Total number of fish</b>		<b>6,539</b>		<b>31,625</b>		<b>2,054</b>		<b>2,717</b>	

### *Offshore waters*

Samples collected from offshore waters in summer and autumn 2022 using gill nets returned 2,768 fish, comprising 24 species (Table 4). This number of fish was similar to that recorded in 2021 (2,933), which was the highest recorded since monitoring began in 2012 and almost 50% more than in 2018 and 2019 (range = 1,125 to 2,933). The 24 species caught in 2022 was also the greatest recorded (range = 17 to 23) and represented almost 70% of all species caught in this monitoring since 2012. As has occurred in most years, the total number of species recorded from each zone in 2022 decreased in an upstream direction from 18 species in the LSCE, to 15 and 13 in the CE and MSE, respectively and 11 in the USE. Values in all zones except the CE were amongst the highest recorded.

As in the ten previous years of monitoring, Perth Herring was among the dominant species in offshore waters overall (55%) and from all four zones, comprising 29–80% of the total catches (Table 4). The Southern Eagle Ray (*Myliobatis tenuicaudata*) and Tailor (*Pomatomus saltatrix*) were abundant in the LSCE (29 and 10% of the catch, respectively). The latter species was also abundant in the CE and MSE, representing 8 and 5% of all fish caught. The Yellowtail Grunter (*Amniataba caudavittata*) was also caught in substantial numbers in the MSE (24%) and USE (38%) as was the Black Bream (*Acanthopagrus butcheri*) in the USE (5%). Catches of several species were relatively high in 2022, including the Southern Eagle Ray, Mulloway (*Argyrosomus japonicus*) and Australian Giant Herring (*Elops machnata*), with values of two, four and five times more than the average between 2012 and 2021, respectively.

**Table 4.** Compositions of the fish communities (CR = Average catch rate [fish/net set] and %C = percentage composition) observed across the six offshore sites sampled in each zone of the Swan Canning Estuary during summer and autumn of 2022. Species ranked by total abundance. Data for the three most abundant species in the catches from each zone are emboldened for emphasis. Species ordered by total abundance throughout the estuary. LSCE = Lower Swan Canning Estuary, CE = Canning Estuary, MSE = Middle Swan Estuary, USE = Upper Swan Estuary.

Species	Common name	LSCE (n = 12)		CE (n = 12)		MSE (n = 12)		USE (n = 12)	
		CR	%C	CR	%C	CR	%C	CR	%C
<i>Nematalosa vlaminghi</i>	Perth Herring	<b>156</b>	<b>29.89</b>	<b>616</b>	<b>79.69</b>	<b>426</b>	<b>62.83</b>	<b>334</b>	<b>42.01</b>
<i>Amniataba caudavittata</i>	Yellowtail Grunter	43	8.24	11	1.42	<b>164</b>	<b>24.19</b>	<b>301</b>	<b>37.86</b>
<i>Pomatomus saltatrix</i>	Tailor	<b>52</b>	<b>9.96</b>	<b>62</b>	<b>8.02</b>	<b>34</b>	<b>5.01</b>	10	1.26
<i>Myliobatis tenuicaudatus</i>	Southern Eagle Ray	<b>153</b>	<b>29.31</b>						
<i>Acanthopagrus butcheri</i>	Black Bream	8	1.53	17	2.20	10	1.47	<b>44</b>	<b>5.53</b>
<i>Mugil cephalus</i>	Sea Mullet			<b>42</b>	<b>5.43</b>	1	0.15	21	2.64
<i>Argyrosomus japonicus</i>	Mulloway			2	0.26	15	2.21	36	4.53
<i>Elops machnata</i>	Australian Giant Herring	3	0.57	2	0.26	8	1.18	36	4.53
<i>Platycephalus westraliae</i>	Yellowtail Flathead	36	6.90	3	0.39	6	0.88	3	0.38
<i>Sillago burrus</i>	West. Trumpeter Whiting	14	2.68			1	0.15		0.00
<i>Rhabdosargus sarba</i>	Tarwhine	8	1.53	7	0.91				
<i>Engraulis australis</i>	Australian Anchovy			2	0.26	4	0.59	8	1.01
<i>Helotes octolineatus</i>	Western Striped Grunter	4	0.77	2	0.26	7	1.03	1	0.13
<i>Gerres subfasciatus</i>	Common Silverbidy	10	1.92	3	0.39	1	0.15		0.00
<i>Arripis georgianus</i>	Australian Herring	9	1.72						
<i>Torquigener pleurogramma</i>	Weeping Toadfish	7	1.34	2	0.26				
<i>Cnidoglanis macrocephalus</i>	Estuary Cobbler	6	1.15	1	0.13				
<i>Sillago schomburgkii</i>	Yellowfin Whiting	5	0.96						
<i>Pseudorhombus jenynsii</i>	Smalltooth Flounder	4	0.77						
<i>Sphyræna obtusata</i>	Yellowtail Barracuda	3	0.57						
<i>Carcharhinus leucas</i>	Bull Shark					1	0.15		0.00
<i>Atherinomorus vaigiensis</i>	Common Hardyhead	1	0.19						
<i>Hippocampus subelongatus</i>	West Australian Seahorse							1	0.13
<i>Pseudocaranx wrighti</i>	Skipjack Trevally			1	0.13				
<b>Total number of species</b>		<b>18</b>		<b>15</b>		<b>13</b>		<b>11</b>	
<b>Average catch rate (fish/net set)</b>		<b>44</b>		<b>64</b>		<b>57</b>		<b>66</b>	
<b>Total number of fish</b>		<b>522</b>		<b>773</b>		<b>678</b>		<b>795</b>	


### 5.3 Ecological condition in 2022

#### Nearshore waters

The ecological condition, based on fish communities, of the nearshore waters of the Swan Canning Estuary was good (B) in summer and good/fair (B/C) in autumn (Fig. 4). The condition of each zone varied substantially during summer (mean FCI scores of 62–72), being best in the USE, MSE and LSCE (good; B) and lowest in the CE with a fair (C) score. By autumn, scores in the CE improved to good, the USE remained in good condition and the LSCE and MSE, declined from good (both zones) to good/fair and fair, respectively (Fig. 4).


Radar plots of the nearshore metric scores for each zone in summer revealed that, in general, zones on the Swan axis of the estuary scored well across all metrics but particularly the *Number of species* (positive metric) and the *Total number of P. olorum*, *Proportion of P. olorum* and *Proportion of detritivores* (all negative metrics; Fig. 5a). High values for the number of species reflect the relatively saline conditions in all zones allowing marine species to occur in areas further upstream than in previous years (see Potter et al., 2016; Valesini et al., 2017). The LSCE scored better in the *Number of trophic generalist species* (negative metric) than the MSE and USE due to large catches of the Western Striped Grunter, a species which spawns in the marine environment and consumes mainly plant material and so is classified as a trophic specialist (Veale et al., 2015; Poh et al., 2018). These fish were fairly small (0+; K. Krispyn, personal observation) and their recruitment could be linked to the strong flows recorded in the preceding year. However, the LSCE scored lower than the MSE and USE for the *Number of estuarine-spawning species*, *Proportion of estuarine-spawning individuals* and *Number of benthic-associated species*. This also occurred in previous years, is due to the presence of ‘marine-like’ salinities in the LSCE which restrict the spatial distribution of estuarine spawning species to areas further upstream.

The fair condition of the CE reflects the lower scores this zone received for the *Proportion of benthic-associated individuals* and the *Total number of P. olorum*. This is driven, in part, by 620 individuals of the Blue Spot Goby caught in the CE, compared to 0, 39 and 149 in the LSCE, MSE and USE, respectively in 2022.




**Figure 4.** Average nearshore Fish Community Index scores and resulting condition grades (A, very good; B, good; C, fair; D, poor; E, very poor) for each zone of the Swan Canning Estuary, and for the estuary as a whole, in summer and autumn of 2022.

(a) Summer 2022



(b) Autumn 2022




**Figure 5.** Average scores (0–10) for each component metric of the nearshore Fish Community Index, calculated from samples collected throughout the LSCE, CE, MSE and USE zones in (a) summer and (b) autumn 2022. Note that an increase in the score for positive metrics (+) reflects an increase in the underlying variable, whereas an increase in the score for negative metrics (-) reflects a decrease in the underlying variable. Therefore, the larger the area covered by the radar plot the better the condition in that zone. Full metric names and explanations are given in Table 1.

In autumn, scores for the CE increased, while those for the LSCE and MSE decreased. Radar plots showed the increase in CE was due mainly to an increase metric scores for the *Number of trophic generalist species* (negative metric) and *Proportion of estuarine-spawning individuals* (positive metric) and to a lesser extent the *Number of trophic specialist species* and *Number of benthic-associated species* (both positive metrics; Fig 5b). It should be noted, however, that the metric scores for the *Proportion of P. olorum* and *Number of species*, both decreased. Trends in some metric scores were consistent across the LSCE, MSE and USE, for example, scores for the *Number of species*, *Number of trophic specialist species* and *Number of benthic-associated species* all declined markedly (i.e. ~3 out

of 10), while the *Proportion of trophic specialists*, *Number of trophic generalist species* increased. In the LSCE, there was also a marked increase in the *Proportion of estuarine-spawning individuals*, which was caused by the absence of the marine-spawning Western Striped Grunter in autumn compared to summer and converse shift in the abundance of the estuarine-spawning Spotted Hardyhead. The *Number of estuarine-spawning species* decreased in both the MSE and USE, reflecting the increasing saline conditions as the salt wedge penetrated further upstream and this was accompanied by a decrease in the Bluespot Goby, increasing scores for the *Proportion of P. olorum* and the *Total number of P. olorum*. Note that there was no increase in these two metrics in the LSCE as the Bluespot Goby was not recorded in this zone in either season and therefore, these metric scores were at their maximum value (10) in both seasons (Fig 5).

### Offshore waters

The ecological condition, based on fish communities, of the offshore waters of the Swan Canning Estuary was good (B) in summer and fair/good (C/B) in autumn (Fig. 6). The condition of each zone varied substantially during summer (mean FCI scores of 49–74), being best in the LSCE and USE (very good; A) and lowest in the CE with a poor (D) score. The MSE received a condition of good (B). By autumn, scores in the LSCE and USE declined from very good to good as did that in the MSE from good to fair/good. While the CE remained in poor condition, its mean FCI score declined from 49 to 45 (Fig. 6).




**Figure 6.** Average offshore Fish Community Index scores and resulting condition grades (A, very good; B, good; C, fair; D, poor; E, very poor) for each zone of the Swan Canning Estuary, and for the estuary as a whole, in summer and autumn of 2022.


Radar plots of the offshore metric scores for the highest scoring zones in summer (LSCE and USE; Fig. 7a) showed very high scores (i.e. > 8 out of 10) for the *Number of species*, *Shannon-Wiener diversity*, *Number of trophic specialists* and the *Proportion of benthic species* (all positive metrics). Values of the first three of these metrics were slightly lower in the MSE, but in excess of those in the CE. The poor condition of the CE was due to very low metric scores for the *Number of species*, *Shannon-Wiener diversity* (both positive metrics) and the *Proportion of detritivores* (negative metric) all typically < 2, albeit very high scores were recorded for the *Proportion of estuarine-spawning individuals* and *Proportion of benthic species*. These trends are due to the fact that, on average, only three species were caught per sample and that > 90% of all fish recorded were Perth Herring, therefore explaining the low measures of diversity and high proportion of detritivores. Hypoxic conditions were

also present through much of the CE during sampling in this season together with the dinoflagellate *Karlodinium* spp. and the raphidophyte *Heterosigma akashiwo* (Appendix vi).

(a) Summer 2022



(b) Autumn 2022




**Figure 7.** Average scores (0–10) for each component metric of the offshore Fish Community Index, calculated from samples collected throughout the LSCE, CE, MSE and USE zones in (a) summer and (b) autumn 2022. Note that an increase in the score for positive metrics (+) reflects an increase in the underlying variable, whereas an increase in the score for negative metrics (-) reflects a decrease in the underlying variable. Therefore, the larger the area covered by the radar plot the better the condition in that zone. Metric names and explanations are given in Table 1.

Whilst the mean offshore FCI scores the LSCE, MSE and USE all declined, they were still greater than that for the CE (Fig. 6). The low scores in the CE reflect very low values for metrics such as the *Proportion of detritivores*, *Number of trophic generalist species* (both negative metrics) and the *Number of trophic specialist species* (positive metric). These were caused by the dominance of detritivores, i.e. Perth Herring and Sea Mullet, which represented 81% of all fish recorded, and the

absence of species such as the Southern Eagle Ray that target benthic invertebrates and are abundant in the adjacent LSCE. The presence of hypoxia in the bottom waters in large extents of the CE during autumn would have deterred benthic species of fish and potentially reduced the availability of their infaunal prey (Tweedley et al., 2016). Although the scores in the LSCE and USE declined, these zones were in good condition and scored well in all metrics except the *Proportion of estuarine-spawning individuals*, due to lower catches of Yellowtail Grunter, particularly in the USE. Condition in the MSE declined to fair/good, with a relatively low metric score for the *Proportion of detritivores* and a decline in the *Number of trophic specialist species* once again reflecting the dominance of Perth Herring. It is relevant, that, although no hypoxia was detected, low concentrations of dissolved oxygen (3-4 mg/L) were recorded in this zone in autumn and may have caused some less tolerant species to move away to more oxic waters.


### Longer term trends in ecological condition

Results indicate that the nearshore waters of the Swan Canning Estuary as a whole were in good condition (B) during 2022, which is better than the relatively consistent overall trend since 2011 of fair/good condition (Fig 8). Aside from 2022, good conditions have only occurred in 2014 and 2016 (Fig. 8). The good scores for 2022 was likely influenced by the strong freshwater flows that occurred in the winter of 2021. Such flows, which were the greatest for 25 years, increased productivity in both the estuary and also nearby coastal waters and facilitated the recruitment of juveniles of large-bodied marine fish species and adults of smaller-bodied and shorter-lived estuaries species.



**Figure 8.** Trend plot of average ( $\pm$ SE) nearshore Fish Community Index (FCI) scores and resulting condition grades (A, very good; B, good; C, fair; D, poor; E, very poor) for the Swan Canning Estuary between 2005 and 2022. Red lines denote boundaries between condition grades.

The mean offshore FCI score for the estuary overall also indicated good condition during 2022, thus the current score of good is in line with the generally upward trend from 2016 onwards (Fig. 9). Moreover, the score in 2022, was only the fourth time good condition has been obtained (i.e. 2012, 2015, 2020 and 2022), with the mean FCI score in 2022 being the highest ever recorded (i.e. 61.4 vs 48.6 to 60.2 in the other 13 years). These good overall scores reflect the consistent very good (A) to fair (C) scores throughout all zones, except for the offshore waters of the CE in summer and autumn (poor; D). It is hypothesised that these good scores in 2022 were driven by the relatively saline and oxic conditions throughout much of the system, particularly in summer, when flows were low. Similarly, good scores in 2012, 2015 and 2020 all coincided with relatively low flow during the January to May period when sampling occurs (i.e. mean = 0.69 ML vs 30.43 GL from 2011-2022). These low levels of flow allow the upstream penetration of saline water facilitating increases in species richness and preventing the occurrence of hypoxia associated with stratification. Conversely, flow was greatest by far in the summer of 2017 (i.e. 282 ML), which had amongst the lowest FCI score due to large spatial extents of hypoxia.



**Figure 9.** Trend plot of average ( $\pm$ SE) offshore Fish Community Index (FCI) scores and resulting condition grades (A, very good; B, good; C, fair; D, poor; E, very poor), for the Swan Canning Estuary as a whole, between 2008 and 2022. Red lines denote boundaries between condition grades.

## Summary

The Fish Community Index (FCI) considers the fish community as a whole and provides an objective means to assess how the structure and function of these communities in shallow, nearshore (< 1.5 m deep) and deeper, offshore waters (> 1.5 m deep) respond to a wide array of stressors affecting the ecosystem. Note that the FCI does not provide information on the population dynamics or health of particular species (in comparison to e.g. Cottingham et al., 2014; Crisp et al., 2018), nor does it provide information on the size or status of the fish stocks in the estuary (e.g. Smith and Lenanton, 2021; Obregón et al., 2022).

Across the entire estuary, the ecological condition of both nearshore and offshore waters in 2022 was assessed as good (B) based on their fish communities (Table 5). Combined, the nearshore and offshore index scores are the highest ever recorded (closely followed by those in 2014), reflecting the fact that the condition in both water depths was good, rather than one good and one fair/good as in 2014 (Hallett and Tweedley, 2014).

The good scores in 2022 were influenced by strong winter flows in most regions combined with the absence of summer rainfall events, relatively saline and oxic conditions and generally low phytoplankton densities through summer and autumn. Along the Swan axis of the estuary the strong freshwater flows that occurred in the winter of 2021 were the greatest for 25 years and will have increased productivity in both the estuary and nearby coastal waters and facilitated the recruitment of fish species. This is evidenced by the number of fish (most of which have an annual lifecycle) recorded in the nearshore waters being by far the largest ever recorded since FCI sampling began.

Overall, the offshore waters of the CE exhibited by far the lowest scores of any zone in 2022. As flows in the Canning River are significantly reduced by regulation (Radin et al., 2007; Norton et al., 2010), this axis of the estuary did not receive the same scale of increases in flow that the USE, MSE and LSCE did. Waters in the CE were once again stratified, hypoxic and, at times, contained moderate levels of the toxic dinoflagellate *Karlodinium* spp.. The poor grades received by the offshore waters of this zone in both seasons are reflective of the trend since the start of regular fish community monitoring in 2012. Over these years the offshore waters of this zone have consistently scored poorly relative to other zones across both seasons, receiving a poor (D) grade in > 50% of monitored seasons. Additional monitoring water quality in this zone has been initiated by DBCA since May 2020 to better understand the factors underlying this trend.

**Table 5.** Fish Community Index (FCI) scores and corresponding ecological condition grades for each zone of the estuary, and the estuary as a whole, during the 2022 monitoring period (mean of all summer and autumn of 2022). LSCE = Lower Swan Canning Estuary, CE = Canning Estuary, MSE = Middle Swan Estuary, USE = Upper Swan Estuary.

	Nearshore		Offshore	
	Mean FCI score	Condition	Mean FCI score	Condition
LSCE	66.02	B	68.74	B
CE	63.64	C/B	46.87	D
MSE	66.70	B	60.94	B
USE	68.98	B	68.76	B
<b>Estuary</b>	<b>66.33</b>	<b>B</b>	<b>61.41</b>	<b>B</b>

## 6. References

- Beatty, S.J., Lear, K.O., Allen, M.G., Lymbery, A.J., Tweedley, J.R., Morgan, D.L., 2022. What factors influence fin-nipping damage by the invasive *Gambusia holbrooki* (Poeciliidae) on native fishes in riverine systems? *Freshw. Biol.* n/a.
- Broadley, A., Stewart-Koster, B., Burford, M.A., Brown, C.J., 2022. A global review of the critical link between river flows and productivity in marine fisheries. *Reviews in Fish Biology and Fisheries*.
- Chuwen, B.M., 2009. Characteristics of the ichthyofaunas of offshore waters in different types of estuary in Western Australia, including the biology of Black Bream *Acanthopagrus butcheri*. Murdoch University, Perth, Australia, p. 213.
- Cottingham, A., Hesp, S.A., Hall, N.G., Hipsey, M.R., Potter, I.C., 2014. Marked deleterious changes in the condition, growth and maturity schedules of *Acanthopagrus butcheri* (Sparidae) in an estuary reflect environmental degradation. *Estuarine, Coastal and Shelf Science* 149, 109-119.
- Crisp, J.A., Loneragan, N.R., Tweedley, J.R., D'Souza, F.M.L., Poh, B., 2018. Environmental factors influencing the reproduction of an estuarine penaeid population and implications for management. *Fisheries Management and Ecology* 25, 203-219.
- Fonseca, V.F., Vasconcelos, R.P., Gamito, R., Pasquaud, S., Gonçalves, C.I., Costa, J.L., Costa, M.J., Cabral, H.N., 2013. Fish community-based measures of estuarine ecological quality and pressure–impact relationships. *Estuarine, Coastal and Shelf Science* 134, 128-137.
- Gibson, G.R., Bowman, M.L., Gerritsen, J., Snyder, B.D., 2000. *Estuarine and Coastal Marine Waters: Bioassessment and Biocriteria Technical Guidance*. Environmental Protection Agency, Office of Water, Washington, DC, EPA 822-B-00-024. U.S.
- Gill, H.S., Potter, I.C., 1993. Spatial segregation amongst goby species within an Australian estuary, with a comparison of the diets and salinity tolerance of the two most abundant species. *Marine Biology* 117, 515-526.
- Gillanders, B.M., Kingsford, M.J., 2002. Impact of changes in flow of freshwater on estuarine and open coastal habitats and the associated organisms. *Oceanography and Marine Biology Annual Review* 40, 233-309.
- Hallett, C.S., 2012. Brief report on fish community responses to the *Karlodinium veneficum* bloom of May 2012, including the response of the Fish Community Index of Estuarine Condition. Murdoch University, Perth, Western Australia, Final report to the Swan River Trust.
- Hallett, C.S., Valesini, F.J., 2012. Validation of the Fish Community Index of Estuarine Condition and development of a monitoring regime for the Swan Canning Riverpark. Murdoch University, Perth, Western Australia.
- Hallett, C.S., Valesini, F.J., Clarke, K.R., Hesp, S.A., Hoeksema, S.D., 2012. Development and validation of fish-based, multimetric indices for assessing the ecological health of Western Australian estuaries. *Estuarine, Coastal and Shelf Science* 104–105, 102-113.
- Hallett, C.S., 2014. Quantile-based grading improves the effectiveness of a multimetric index as a tool for communicating estuarine condition. *Ecological Indicators* 39, 84-87.
- Hallett, C.S., Tweedley, J.R., 2014. Assessment of the condition of the Swan Canning Estuary in 2014, based on the Fish Community Indices of estuarine condition. Murdoch University, Perth, Western Australia, Final report to the Swan River Trust, p. 32.

- Hallett, C.S., Valesini, F.J., Clarke, K.R., Hoeksema, S.D., 2016. Effects of a harmful algal bloom on the community ecology, movements and spatial distributions of fishes in a microtidal estuary. *Hydrobiologia* 763, 267-284.
- Hallett, C.S., 2017. Assessment of the condition of the Swan Canning Estuary in 2017, based on the Fish Community Indices of estuarine condition. Murdoch University, Perth, Western Australia, Final report to the Department of Biodiversity, Conservation and Attractions, p. 41.
- Hallett, C.S., Hobday, A.J., Tweedley, J.R., Thompson, P.A., McMahon, K., Valesini, F.J., 2018. Observed and predicted impacts of climate change on the estuaries of south-western Australia, a Mediterranean climate region. *Regional Environmental Change* 18, 1357-1373.
- Hallett, C.S., Trayler, K.M., Valesini, F.J., 2019. The Fish Community Index: A practical management tool for monitoring and reporting estuarine ecological condition. *Integrated Environmental Assessment and Management* 15, 726-738.
- Hodgkin, E.P., Hesp, P., 1998. Estuaries to salt lakes: Holocene transformation of the estuarine ecosystems of south-western Australia. *Marine and Freshwater Research* 49, 183-201.
- Hoeksema, S.D., Chuwen, B.M., Potter, I.C., 2009. Comparisons between the characteristics of ichthyofaunas in nearshore waters of five estuaries with varying degrees of connectivity with the ocean. *Estuarine, Coastal and Shelf Science* 85, 22-35.
- Hogan-West, K., Tweedley, J.R., Coulson, P.G., Poh, B., Loneragan, N.R., 2019. Abundance and distribution of the non-indigenous *Acentrogobius pflaumii* and native gobiids in a temperate Australian estuary. *Estuaries and Coasts* 42, 1612-1631.
- Krispyn, K.N., Loneragan, N.R., Whitfield, A.K., Tweedley, J.R., 2021. Salted mullet: Protracted occurrence of *Mugil cephalus* under extreme hypersaline conditions. *Estuarine, Coastal and Shelf Science* 261, 107533.
- Norton, S., Storer, T., Galvin, L., 2010. Ecological study of the lower Canning River environmental water releases. Department of Water, Government of Western Australia, Perth, p. 43.
- Obregón, C., Christensen, J., Zeller, D., Hughes, M., Tweedley, J.R., Gaynor, A., Loneragan, N.R., 2022. Local fisher knowledge reveals changes in size of blue swimmer crabs in small-scale fisheries. *Marine Policy*.
- Poh, B., Tweedley, J.R., Chaplin, J.A., Trayler, K.M., Loneragan, N.R., 2018. Estimating predation rates of restocked individuals: The influence of timing-of-release on metapenaeid survival. *Fisheries Research* 198, 165-179.
- Potter, I.C., Tweedley, J.R., Elliott, M., Whitfield, A.K., 2015a. The ways in which fish use estuaries: A refinement and expansion of the guild approach. *Fish and Fisheries* 16, 230-239.
- Potter, I.C., Warwick, R.M., Hall, N.G., Tweedley, J.R., 2015b. The physico-chemical characteristics, biota and fisheries of estuaries, in: Craig, J. (Ed.), *Freshwater Fisheries Ecology*. Wiley-Blackwell, pp. 48-79.
- Potter, I.C., Veale, L.J., Tweedley, J.R., Clarke, K.R., 2016. Decadal changes in the ichthyofauna of a eutrophic estuary following a remedial engineering modification and subsequent environmental shifts. *Estuarine, Coastal and Shelf Science* 181, 345-363.

- Prince, J., Potter, I., 1983. Life-cycle duration, growth and spawning times of five species of atherinidae (Teleostei) found in a Western Australian estuary. *Marine and Freshwater Research* 34, 287-301.
- Radin, L., La Spina, K., Braimbridge, M., Malseed, B., 2007. Environmental values, flow related issues and objectives for the Canning River, Western Australia. Department of Water, Government of Western Australia, Perth, p. 90.
- Smith, K.A., Lenanton, R.C.J., 2021. Almost forgotten: Historical abundance of eel-tail catfish populations in south-western Australian estuaries and their decline due to habitat loss and historical overfishing. *Regional Studies in Marine Science* 41, 101605.
- Tweedley, J.R., Hallett, C.S., Warwick, R.M., Clarke, K.R., Potter, I.C., 2016. The hypoxia that developed in a microtidal estuary following an extreme storm produced dramatic changes in the benthos. *Marine and Freshwater Research* 67, 327-341.
- Tweedley, J.R., Warwick, R.M., Hallett, C.S., Potter, I.C., 2017. Fish-based indicators of estuarine condition that do not require reference data. *Estuarine, Coastal and Shelf Science* 191, 209-220.
- Tweedley, J.R., Krispyn, K.N., Hallett, C.S., 2021. Swan Canning Estuary condition assessment based on fish communities - 2020. Murdoch University, Perth, Western Australia, Final report to the Department of Biodiversity, Conservation and Attractions, p. 47.
- Valesini, F.J., Coen, N.J., Wildsmith, M.D., Hourston, M., Tweedley, J.R., Hallett, C.S., Linke, T.E., Potter, I.C., 2009. Relationships between fish faunas and habitat type in south-western Australian estuaries. Project 2004/045. Draft Final Report for Fisheries Research and Development Corporation. Murdoch University, Perth.
- Valesini, F.J., Hallett, C.S., Cottingham, A., Hesp, S.A., Hoeksema, S.D., Hall, N.G., Linke, T.E., Buckland, A.J., 2011. Development of biotic indices for establishing and monitoring ecosystem health of the Swan Canning Estuary. Murdoch University, Perth, Western Australia, Final Report to the Swan River Trust, Department of Water, Department of Fisheries.
- Valesini, F.J., Cottingham, A., Hallett, C.S., Clarke, K.R., 2017. Interdecadal changes in the community, population and individual levels of the fish fauna of an extensively modified estuary. *Journal of Fish Biology* 90, 1734-1767.
- Veale, L., Tweedley, J.R., Clarke, K.R., Hallett, C.S., Potter, I.C., 2014. Characteristics of the ichthyofauna of a temperate microtidal estuary with a reverse salinity gradient, including inter-decadal comparisons. *Journal of Fish Biology* 85, 1320-1354.
- Veale, L., Coulson, P., Hall, N., Hesp, A., Potter, I.C., 2015. Age and size compositions, habitats, growth and reproductive characteristics of a terapontid (*Pelates octolineatus*) in coastal waters. *Marine and Freshwater Research* 66, 535-548.
- Villéger, S., Miranda, J.R., Hernández, D.F., Mouillot, D., 2010. Contrasting changes in taxonomic vs. functional diversity of tropical fish communities after habitat degradation. *Ecological Applications* 20, 1512-1522.
- Whitfield, A.K., Elliott, M., 2002. Fishes as indicators of environmental and ecological changes within estuaries: a review of progress and some suggestions for the future. *Journal of Fish Biology* 61, 229-250.

Whitfield, A.K., Elliott, M., Blaber, S.J.M., Harrison, T.D., Tweedley, J.R., Franco, A., Potter, I.C., 2022. Fish Assemblages and Functional Groups, in: Whitfield, A.K., Elliott, M., Blaber, S.J.M., Able, K. (Eds.), Estuarine Fish and Fisheries. Wiley.

## 7. Appendices

**Appendix (i).** Descriptions of (a) nearshore and (b) offshore Fish Community Index monitoring sites. LSCE, Lower Swan Canning Estuary; CE, Canning Estuary; MSE, Middle Swan Estuary; USE, Upper Swan Estuary.

Zone	Site Code	Lat-Long (S, E)	Description	
<b>(a) – Nearshore</b>				
LSCE	LSCE3	-32°01'29", 115°46'27"	Shoreline in front of vegetation on eastern side of Point Roe, Mosman Pk	
	LSCE4	-31°59'26", 115°47'08"	Grassy shore in front of houses to east of Claremont Jetty	
	LSCE5	-32°00'24", 115°46'52"	North side of Point Walter sandbar	
	LSCE6	-32°01'06", 115°48'19"	Shore in front of bench on Attadale Reserve	
	LSCE7	-32°00'11", 115°50'29"	Sandy bay below Point Heathcote	
	LSCE8	-31°59'11", 115°49'40"	Eastern side of Pelican Point, immediately south of sailing club	
	CE	CE1	-32°01'28", 115°51'16"	Sandy shore to south of Deepwater Point boat ramp
		CE2	-32°01'54", 115°51'33"	Sandy beach immediately to north of Mount Henry Bridge
CE5		-32°01'40", 115°52'58"	Bay in Shelley Beach, adjacent to jetty	
CE6		-32°01'29", 115°53'11"	Small clearing in vegetation off North Riverton Drive	
CE7		-32°01'18", 115°53'43"	Sandy bay in front of bench, east of Wadjup Point	
CE8		-32°01'16", 115°55'14"	Sandy beach immediately downstream of Kent Street Weir	
MSE		MSE2	-31°58'12", 115°51'07"	Sandy beach on South Perth foreshore, west of Mends St Jetty
		MSE4	-31°56'34", 115°53'06"	Shoreline in front of Belmont racecourse, north of Windan Bridge
	MSE5	-31°56'13", 115°53'23"	Beach to west of jetty in front of Maylands Yacht Club	
	MSE6	-31°57'13", 115°53'56"	Small beach upstream of Belmont Water Ski Area boat ramp	
	MSE7	-31°55'53", 115°55'10"	Beach in front of scout hut, east of Garratt Road Bridge	
	MSE8	-31°55'37", 115°56'18"	Vegetated shoreline, Claughton Reserve, upstream of boat ramp	
	USE	USE1	-31°55'20", 115°57'03"	Small beach adjacent to jetty at Sandy Beach Reserve, Bassendean
		USE3	-31°53'43", 115°57'32"	Sandy bay opposite Bennett Brook, at Fishmarket Reserve, Guildford
USE4		-31°53'28", 115°58'32"	Shoreline in front of Guildford Grammar stables, opposite Lilac Hill Park	
USE5		-31°53'13", 115°59'29"	Small, rocky beach after bend in river at Ray Marshall Park	
USE6		-31°52'41", 115°59'31"	Small beach with iron fence, in front of Caversham house	
USE7		-31°52'22", 115°59'39"	Sandy shore on bend in river, below house on hill, upstream of powerlines	
<b>(b) – Offshore</b>				
LSCE	LSCE1G	-32°00'24", 115°46'56"	In deeper water <i>ca</i> 100 m off north side of Point Walter sandbar	
	LSCE2G	-32°00'12", 115°48'07"	Alongside seawall west of Armstrong Spit, Dalkeith	
	LSCE3G	-32°01'00", 115°48'44"	Parallel to shoreline, running westwards from Beacon 45, Attadale	
	LSCE4G	-32°00'18", 115°50'01"	In deep water of Waylen Bay, from <i>ca</i> 50 m east of Applecross jetty	
	LSCE5G	-31°59'37", 115°51'09"	Perpendicular to Como Jetty, running northwards	
	LSCE6G	-31°59'12", 115°49'42"	<i>Ca</i> 20 m from, and parallel to, sandy shore on east side of Pelican Point	
CE	CE1G	-32°01'58", 115°51'36"	Underneath Mount Henry Bridge, parallel to northern shoreline	
	CE2G	-32°01'48", 115°51'46"	Parallel to, and <i>ca</i> 20 m from, western shoreline of Aquinas Bay	
	CE3G	-32°01'49", 115°52'19"	To north of navigation markers, Aquinas Bay	
	CE4G	-32°01'48", 115°52'33"	Adjacent to Old Post Line (SW-ern end; Salter Point)	
	CE5G	-32°01'36", 115°52'52"	Adjacent to Old Post Line (NE-ern end; Prisoner Point)	
	CE6G	-32°01'20", 115°53'15"	Adjacent to Old Post Line, Shelley Water	
MSE	MSE1G	-31°58'03", 115°51'03"	From jetty at Point Belches towards Mends St Jetty, Perth Water	
	MSE2G	-31°56'57", 115°53'05"	Downstream of Windan Bridge, parallel to Burswood shoreline	
	MSE3G	-31°56'22", 115°53'05"	Downstream from port marker, parallel to Joel Terrace, Maylands	
	MSE4G	-31°57'13", 115°54'12"	Parallel to shore from former boat shed jetty, Cracknell Park, Belmont	
	MSE5G	-31°55'57", 115°55'12"	Parallel to southern shoreline, upstream of Garratt Road Bridge	
	MSE6G	-31°55'23", 115°56'25"	Parallel to eastern bank at Garvey Pk, from south of Ron Courtney Island	
USE	USE1G	-31°55'19", 115°57'09"	Parallel to tree-lined eastern bank, upstream of Sandy Beach Reserve	
	USE2G	-31°53'42", 115°57'40"	Along northern riverbank, running upstream from Bennett Brook	
	USE3G	-31°53'16", 115°58'42"	Along northern bank on bend in river, to north of Lilac Hill Park	
	USE4G	-31°53'17", 115°59'23"	Along southern bank, downstream from bend at Ray Marshall Pk	
	USE5G	-31°52'13", 115°59'40"	Running along northern bank, upstream from Sandalford winery jetty	
	USE6G	-31°52'13", 116°00'18"	Along southern shore adjacent to Midland Brickworks, from outflow pipe	

## **Appendix (ii).** Descriptions of sampling and processing procedures.

### ***Nearshore sampling methods***

- On each sampling occasion, one replicate sample of the nearshore fish community is collected from each of the fixed, nearshore sampling sites.
- Sampling is not conducted during or within 3-5 days following any significant flow event.
- Nearshore fish samples are collected using a beach seine net that is 21.5 m long, comprises two 10 m-long wings (6 m of 9 mm mesh and 4 m of 3 mm mesh) and a 1.5 m-long bunt (3 mm mesh) and fishes to a depth of 1.5 m.
- This net is walked out from the beach to a maximum depth of approximately 1.5 m and deployed parallel to the shore, and is then rapidly dragged towards and onto the shore, so that it sweeps a roughly semicircular area of approximately 116 m<sup>2</sup>.
- If a seine net deployment returns a catch of fewer than five fish, an additional sample is performed at the site (separated from the first sample by either 15 minutes or by 10-20 m distance). In the event that more than five fish are caught in the second sample, this second replicate is then used as the sample for that site and those fish from the first sample returned to the water alive. If, however, 0-5 fish are again caught, the original sample can be assumed to have been representative of the fish community present and be used as the sample for that site. The fish from the latter sample are then returned alive to the water. The above procedure thus helps to identify whether a collected sample is representative of the fish community present and enables instances of false negative catches to be identified and eliminated.
- Once an appropriate sample has been collected, any fish that may be readily identified to species (e.g. those larger species which are caught in relatively lower numbers) are counted and returned to the water alive.
- All other fish caught in the nets are placed into zip-lock polythene bags, euthanised in an ice slurry and preserved on ice in eskies in the field, except in cases where large catches (e.g. thousands) of small fish are obtained. In such cases, an appropriate sub-sample (e.g. one half to one eighth of the entire catch) is retained and the remaining fish are returned alive to the water. All retained fish are then bagged and frozen until their identification in the laboratory.

### ***Offshore sampling methods***

- On each sampling occasion, one replicate sample of the offshore fish community is collected from each of the fixed, offshore sampling sites.
- Sampling is not conducted within 3-5 days following any significant flow event.
- Offshore fish samples are collected using a sunken, multimesh gill net that consists of eight 20 m-long panels with stretched mesh sizes of 35, 51, 63, 76, 89, 102, 115 and 127 mm. These nets are deployed (i.e. laid parallel to the bank) from a boat immediately before sunset and retrieved after three hours.
- Given the time and labour associated with offshore sampling and the need to monitor the set nets for safety purposes, a maximum of three replicate net deployments is performed within a single zone in any one night. The three nets are deployed sequentially, and retrieved in the same order.
- During net retrieval (and, typically, when catch rates are sufficiently low to allow fish to be removed rapidly in the course of retrieval), any fishes that may be removed easily from the net are carefully removed, identified, counted, recorded and returned to the water alive as the net is pulled into the boat.

- All other fish caught in the nets are removed once the net has been retrieved. Retained fish are placed into zip-lock polythene bags in an ice slurry, preserved on ice in eskies in the field, and subsequently frozen until their identification in the laboratory.

Following their identification to the lowest possible taxon in the field or laboratory by fish specialists trained in fish taxonomy, all assigned scientific and common names are checked and standardised by referencing the Checklist of Australian Aquatic Biota (CAAB) database (Rees *et al.* on-line version), and the appropriate CAAB species code is allocated to each species. The abundance data for each species in each sample is entered into a database for record and subsequent computation of the biotic indices.

Rees, A.J.J., Yearsley, G.K., Gowlett-Holmes, K. and Pogonoski, J. Codes for Australian Aquatic Biota (on-line version). CSIRO Marine and Atmospheric Research, World Wide Web electronic publication, 1999 onwards. Available at: <http://www.cmar.csiro.au/caab/>. Last accessed 29 January 2021.


**Appendix (iii).** List of species caught from the Swan Canning Estuary, and their functional guilds: D, Demersal; P, Pelagic; BP, Benthopelagic; SP, Small pelagic; SB, Small benthic; MS, Marine straggler; MM, Marine migrant; SA, Semi-anadromous; ES, Estuarine species; FM, Freshwater migrant; ZB, Zoobenthivore; PV, Piscivore; ZP, Zooplanktivore; DV, Detritivore; OV, Omnivore/Opportunist; HV, Herbivore. See Potter et al. (2015a); Whitfield et al. (2022) for descriptions of the guilds.

Species name	Common name	Family	Habitat Guild	Estuarine Use Guild	Feeding Mode Guild
<i>Heterodontus portusjacksoni</i>	Port Jackson Shark	Heterodontidae	D	MS	ZB
<i>Carcharhinus leucas</i>	Bull Shark	Carcharhinidae	P	MS	PV
<i>Myliobatis tenuicaudatus</i>	Southern Eagle Ray	Myliobatidae	D	MS	ZB
<i>Elops machnata</i>	Australian Giant Herring	Elopidae	BP	MS	PV
<i>Sardinops sagax</i>	Australian Sardine	Clupeidae	P	MS	ZP
<i>Spratelloides robustus</i>	Blue Sprat	Clupeidae	SP	MM	ZP
<i>Hyperlophus vittatus</i>	Sandy Sprat	Clupeidae	SP	MM	ZP
<i>Nematalosa vlaminghi</i>	Perth Herring	Clupeidae	BP	SA	DV
<i>Sardinella lemuru</i>	Scaly Mackerel	Clupeidae	P	MS	ZP
<i>Engraulis australis</i>	Australian Anchovy	Engraulidae	SP	ES	ZP
<i>Galaxias occidentalis</i>	Western Galaxias	Galaxiidae	SB	FM	ZB
<i>Carassius auratus</i>	Goldfish	Cyprinidae	BP	FM	OV
<i>Cnidoglanis macrocephalus</i>	Estuary Cobbler	Plotosidae	D	MM	ZB
<i>Tandanus bostocki</i>	Freshwater Cobbler	Plotosidae	D	FM	ZB
<i>Hyporhamphus melanochir</i>	Southern Garfish	Hemiramphidae	P	ES	HV
<i>Hyporhamphus regularis</i>	River Garfish	Hemiramphidae	P	FM	HV
<i>Gambusia holbrooki</i>	Eastern Gambusia	Poeciliidae	SP	FM	ZB
<i>Leptatherina presbyteroides</i>	Silver Fish	Atherinidae	SP	MM	ZP
<i>Atherinomorus vaigiensis</i>	Common Hardyhead	Atherinidae	SP	MM	ZB
<i>Atherinosoma elongatum</i>	Elongate Hardyhead	Atherinidae	SP	ES	ZB
<i>Leptatherina wallacei</i>	Western Hardyhead	Atherinidae	SP	ES	ZP
<i>Craterocephalus mugiloides</i>	Spotted Hardyhead	Atherinidae	SP	ES	ZB
<i>Cleidopus gloriamaris</i>	Australian Pineapplefish	Monocentrididae	D	MS	ZB
<i>Phyllopteryx taeniolatus</i>	Common Seadragon	Syngnathidae	D	MS	ZB
<i>Hippocampus angustus</i>	Western Spiny Seahorse	Syngnathidae	D	MS	ZP
<i>Urocampus carinirostris</i>	Hairy Pipefish	Syngnathidae	D	ES	ZP
<i>Stigmatopora argus</i>	Spotted Pipefish	Syngnathidae	D	MS	ZP
<i>Stigmatopora nigra</i>	Widebody Pipefish	Syngnathidae	D	MS	ZB
<i>Pugnaso curtirostris</i>	Pugnose Pipefish	Syngnathidae	D	MS	ZP
<i>Vanacampus phillipi</i>	Port Phillip Pipefish	Syngnathidae	D	MS	ZB
<i>Filicampus tigris</i>	Tiger Pipefish	Syngnathidae	D	MS	ZP
<i>Gymnapistes marmoratus</i>	Soldier	Tetrarogidae	D	MS	ZB
<i>Chelidonichthys kumu</i>	Red Gurnard	Triglidae	D	MS	ZB
<i>Leviprora inops</i>	Longhead Flathead	Platycephalidae	D	MS	PV
<i>Platycephalus laevigatus</i>	Rock Flathead	Platycephalidae	D	MS	PV
<i>Platycephalus westraliae</i>	Yellowtail Flathead	Platycephalidae	D	ES	PV
<i>Pegasus lancifer</i>	Sculptured Seamothe	Pegasidae	D	MS	ZB
<i>Nannoperca vittata</i>	Western Pygmy Perch	Percichthyidae	BP	FM	ZB
<i>Amniataba caudavittata</i>	Yellowtail Grunter	Terapontidae	BP	ES	OP
<i>Bidyanus bidyanus</i>	Silver Perch	Terapontidae	BP	FM	OV
<i>Helotes octolineatus</i>	Western Striped Grunter	Terapontidae	BP	MM	OV

Species name	Common name	Family	Habitat Guild	Estuarine Use Guild	Feeding Mode Guild
<i>Pelsartia humeralis</i>	Sea Trumpeter	Terapontidae	BP	MS	OV
<i>Siphamia cephalotes</i>	Wood's Siphonfish	Apogonidae	BP	MS	ZB
<i>Ostorhinchus rueppellii</i>	Western Gobbleguts	Apogonidae	BP	ES	ZB
<i>Sillaginodes punctatus</i>	King George Whiting	Sillaginidae	D	MM	ZB
<i>Sillago bassensis</i>	Southern School Whiting	Sillaginidae	D	MS	ZB
<i>Sillago burrus</i>	Western Trumpeter Whiting	Sillaginidae	D	MM	ZB
<i>Sillago schomburgkii</i>	Yellowfin Whiting	Sillaginidae	D	MM	ZB
<i>Sillago vittata</i>	Western School Whiting	Sillaginidae	D	MM	ZB
<i>Pomatomus saltatrix</i>	Tailor	Pomatomidae	P	MM	PV
<i>Trachurus novaezelandiae</i>	Yellowtail Scad	Carangidae	P	MS	ZB
<i>Scomberoides tol</i>	Needleskin Queenfish	Carangidae	P	MS	PV
<i>Pseudocaranx georgianus</i>	Silver Trevally	Carangidae	BP	MM	ZB
<i>Pseudocaranx wrighti</i>	Skipjack Trevally	Carangidae	BP	MM	ZB
<i>Arripis georgianus</i>	Australian Herring	Arripidae	P	MM	PV
<i>Pentapodus vitta</i>	Western Butterfish	Nemipteridae	BP	MS	ZB
<i>Gerres subfasciatus</i>	Common Silverbiddy	Gerreidae	BP	MM	ZB
<i>Acanthopagrus butcheri</i>	Black Bream	Sparidae	BP	ES	OP
<i>Rhabdosargus sarba</i>	Tarwhine	Sparidae	BP	MM	ZB
<i>Argyrosomus japonicus</i>	Mulloway	Sciaenidae	BP	MM	PV
<i>Parupeneus spilurus</i>	Blacksaddle Goatfish	Mullidae	D	MS	ZB
<i>Neatypus obliquus</i>	Footballer Sweep	Scorpididae	P	MS	ZP
<i>Scorpius aequipinnis</i>	Sea Sweep	Scorpididae	P	MS	ZP
<i>Enoplosus armatus</i>	Old Wife	Enoplosidae	D	MS	ZB
<i>Geophagus brasiliensis</i>	[a cichlid]	Cichlidae	BP	FM	OV
<i>Aldrichetta forsteri</i>	Yelloweye Mullet	Mugilidae	P	MM	OV
<i>Mugil cephalus</i>	Sea Mullet	Mugilidae	P	MM	DV
<i>Sphyraena novaehollandiae</i>	Snook	Sphyraenidae	P	MS	PV
<i>Sphyraena obtusata</i>	Striped Barracuda	Sphyraenidae	P	MS	PV
<i>Neodax balteatus</i>	Little Weed Whiting	Labridae	D	MS	OV
<i>Siphonognathus radiatus</i>	Longray Weed Whiting	Labridae	D	MS	OV
<i>Haletta semifasciata</i>	Blue Weed Whiting	Labridae	D	MS	OV
<i>Heteroscarus acroptilus</i>	Rainbow Cale	Labridae	D	MS	OV
<i>Parapercis haackei</i>	Wavy Grubfish	Pinguipedidae	D	MS	ZB
<i>Lesueurina platycephala</i>	Flathead Sandfish	Leptoscopidae	D	MS	ZB
<i>Istiblennius meleagris</i>	Peacock Rockskipper	Blenniidae	D	MS	HV
<i>Omobranchus germaini</i>	Germain's Blenny	Blenniidae	SB	MS	ZB
<i>Parablennius intermedius</i>	Horned Blenny	Blenniidae	D	MS	ZB
<i>Parablennius postocolomaculatus</i>	False Tasmanian Blenny	Blenniidae	SB	MS	OV
<i>Petroscirtes breviceps</i>	Shorthead Sabretooth Blenny	Blenniidae	SB	MS	OV
<i>Cristiceps australis</i>	Southern Crested Weedfish	Clinidae	D	MS	ZB
<i>Pseudocalliurichthys goodladi</i>	Longspine Dragonet	Callionymidae	D	MS	ZB
<i>Eocallionymus papilio</i>	Painted Stinkfish	Callionymidae	D	MS	ZB
<i>Callogobius mucosus</i>	Sculptured Goby	Gobiidae	SB	MS	ZB
<i>Favonigobius lateralis</i>	Southern Longfin Goby	Gobiidae	SB	MM	ZB
<i>Nesogobius pulchellus</i>	Sailfin Goby	Gobiidae	SB	MS	ZB
<i>Arenigobius bifrenatus</i>	Bridled Goby	Gobiidae	SB	ES	ZB


Species name	Common name	Family	Habitat Guild	Estuarine Use Guild	Feeding Mode Guild
<i>Pseudogobius olorum</i>	Bluespot Goby	Gobiidae	SB	ES	OV
<i>Bathygobius krefftii</i>	Krefft's Frillgoby	Gobiidae	SB	MM	ZB
<i>Callogobius depressus</i>	Flathead Goby	Gobiidae	SB	MS	ZB
<i>Favonigobius punctatus</i>	Yellowspotted Sandgoby	Gobiidae	SB	ES	ZB
<i>Afurcagobius suppositus</i>	Southwestern Goby	Gobiidae	SB	ES	ZB
<i>Redigobius macrostoma</i>	Largemouth Goby	Gobiidae	SB	ES	ZB
<i>Tridentiger trignocephalus</i>	Trident Goby	Gobiidae	SB	MS	ZB
<i>Pseudorhombus jenynsii</i>	Smalltooth Flounder	Paralichthyidae	D	MM	ZB
<i>Ammotretis rostratus</i>	Longsnout Flounder	Pleuronectidae	D	MM	ZB
<i>Ammotretis elongatus</i>	Elongate Flounder	Pleuronectidae	D	MM	ZB
<i>Cynoglossus broadhursti</i>	Southern Tongue Sole	Cynoglossidae	D	MS	ZB
<i>Acanthaluteres brownii</i>	Spinytail Leatherjacket	Monacanthidae	D	MS	OV
<i>Acanthaluteres vittiger</i>	Toothbrush Leatherjacket	Monacanthidae	D	MS	OV
<i>Eubalichthys mosaicus</i>	Mosaic Leatherjacket	Monacanthidae	D	MS	OV
<i>Scobinichthys granulatus</i>	Rough Leatherjacket	Monacanthidae	D	MS	OV
<i>Monacanthus chinensis</i>	Fanbelly Leatherjacket	Monacanthidae	D	MM	OV
<i>Chaetodermis penicilligerus</i>	Tasselled Leatherjacket	Monacanthidae	D	MS	OV
<i>Brachaluteres jacksonianus</i>	Southern Pygmy Leatherjacket	Monacanthidae	D	MS	OV
<i>Meuschenia freycineti</i>	Sixspine Leatherjacket	Monacanthidae	D	MM	OV
<i>Acanthaluteres spilomelanurus</i>	Bridled Leatherjacket	Monacanthidae	D	MM	OV
<i>Torquigener pleurogramma</i>	Weeping Toadfish	Tetraodontidae	BP	MM	OP
<i>Contusus brevicaudus</i>	Prickly Toadfish	Tetraodontidae	BP	MS	OP
<i>Polyspina piosae</i>	Orangebarred Puffer	Tetraodontidae	BP	MS	OP
<i>Diodon nictemerus</i>	Globefish	Diodontidae	D	MS	ZB

**Appendix (iv).** (a) Total annual flow between 1971 and 2021 and (b) total monthly flow between January 2011 and May 2022 at Walyunga on the Swan River (gauging station 16401). Data from 2021 highlighted in black in (a) and as a dashed line in (b). Data recorded by the Department of Water and Environmental Regulation and extracted from <https://wir.water.wa.gov.au/Pages/Water-Information-Reporting.aspx>.




**Appendix (v).** (a) Total annual flow between 1971 and 2021 and (b) total monthly flow between January 2011 and May 2022 at Seaforth on the Canning River (gauging station 16417). Data from 2021 highlighted in black in (a) and as a dashed line in (b). Data recorded by the Department of Water and Environmental Regulation and extracted from <https://wir.water.wa.gov.au/Pages/Water-Information-Reporting.aspx>.


(a)



(b)




**Appendix (vi).** A representative selection of vertical contour plots of salinity, dissolved oxygen concentrations (mg/L), Chlorophyll fluorescence ( $\mu\text{g/L}$ ) and water temperature ( $^{\circ}\text{C}$ ) measured at monitoring stations along the length of the Swan Canning Estuary on occasions throughout the summer to autumn period of fish community sampling. Prepared by the Department of Biodiversity, Conservation and Attractions (<https://www.dbca.wa.gov.au/science/riverpark-monitoring>).




# LSCE, MSE and USE zones in summer through autumn 2022


4<sup>th</sup> January 2022




10<sup>th</sup> January 2022




17<sup>th</sup> January 2022




24<sup>th</sup> January 2022




31<sup>st</sup> January 2022




7<sup>th</sup> February 2022




14<sup>th</sup> February 2022




21<sup>st</sup> February 2022




28<sup>th</sup> February 2022




8<sup>th</sup> March 2022




14<sup>th</sup> March 2022




21<sup>st</sup> March 2022




28<sup>th</sup> March 2022




4<sup>th</sup> April 2022




11<sup>th</sup> April 2022




19<sup>th</sup> April 2021




26<sup>th</sup> April 2021



2<sup>nd</sup> May 2021




9th May 2021




# CE zone in summer through autumn 2022


5<sup>th</sup> January 2022




11<sup>th</sup> January 2022




18<sup>th</sup> January 2022




25<sup>th</sup> January 2022




1<sup>st</sup> February 2022




8<sup>th</sup> February 2022




15<sup>th</sup> February 2022




22<sup>nd</sup> February 2022




1<sup>st</sup> March 2022




9<sup>th</sup> March 2022




15<sup>th</sup> March 2022




22<sup>nd</sup> March 2022




29<sup>th</sup> March 2022




5<sup>th</sup> April 2022




12<sup>th</sup> April 2022




20<sup>th</sup> April 2021



27<sup>th</sup> April 2021



3<sup>rd</sup> May 2021



10<sup>th</sup> May 2021

