



# 2025 Commercial Kangaroo Harvest Quota Submission for Western Australia

For submission under the Commonwealth *Environment  
Protection and Biodiversity Conservation Act 1999*.

January 2025

Department of Biodiversity, Conservation and Attractions  
Locked Bag 104  
Bentley Delivery Centre WA 6983  
Phone: (08) 9219 9000  
Fax: (08) 9334 0498

[www.dbca.wa.gov.au](http://www.dbca.wa.gov.au)

© Department of Biodiversity, Conservation and Attractions on behalf of the State of Western Australia 2025  
January 2025

This work is copyright. You may download, display, print and reproduce this material in unaltered form (retaining this notice) for your personal, non-commercial use or use within your organisation. Apart from any use as permitted under the *Copyright Act 1968*, all other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to the Department of Biodiversity, Conservation and Attractions.

Questions regarding the use of this material should be directed to:  
Senior Conservation Zoologist  
Species and Communities Program  
Department of Biodiversity, Conservation and Attractions  
Locked Bag 104  
Bentley Delivery Centre WA 6983  
Email: [fauna.recovery@dbca.wa.gov.au](mailto:fauna.recovery@dbca.wa.gov.au)

The recommended reference for this publication is:  
Department of Biodiversity, Conservation and Attractions, 2025, *2025 Commercial Kangaroo Harvest Quota Submission for Western Australia*, Department of Biodiversity, Conservation and Attractions, Perth.

This document is available in alternative formats on request.

# Contents

1	Proposed quotas for Western Australia for 2025 .....	5
1.1	State summary .....	7
1.2	Regional quotas for red kangaroos in 2025 .....	7
1.3	Regional quotas for western grey kangaroos in 2025 .....	8
1.4	South-East Kangaroo Management Area quotas for western grey kangaroos in 2025 .....	8
2	Aerial survey and population estimation methods .....	11
2.1	Fixed-wing aerial survey .....	11
2.2	Population estimates .....	12
3	Criteria used in setting quotas .....	12
4	Harvest monitoring .....	13
	References .....	14
	Appendices .....	15

## Appendices

Appendix 1	Harvest monitoring results for Western Australia .....	15
Appendix 2	Rainfall and drought maps .....	19
Appendix 3	Regional population estimates .....	22
	Northern Zone .....	22
	Central Zone .....	26
	South-East Zone .....	31
	South-West Zone .....	36

## Figures

Figure 1:	Kangaroo Population Monitoring Zones over the allowable harvest areas in Western Australia .....	9
Figure 2:	Kangaroo Management Areas in WA .....	10

## Tables

Table 1	Aerial survey parameters .....	11
Table 2:	Standard fixed-wing survey correction factors used for population estimates. ....	12
Table 3:	Density thresholds for Western Australia PMZ for red and western grey kangaroos .....	13



# 1 Proposed quotas for Western Australia for 2025

This document presents the proposed quotas for commercial kangaroo harvest in Western Australia for 2025. Quotas are set for pre-determined Population Management Zones (PMZ) or Kangaroo Management Areas (KMA) (Figure 1 and Figure 2). This document should be read and considered in conjunction with the *Management Plan for the Commercial Harvest of Kangaroos in Western Australia 2024-2028* (Management Plan).

Scheduled aerial surveys for the Central PMZ and South-West PMZ did not occur in 2024. Population estimates in 2024 for all PMZ or KMA were calculated using the most recent population estimate adjusted for regional rainfall and commercial harvest offtake, as per the requirements of the Management Plan.

Western Australia experienced unusually dry and hot summer weather across multiple areas of the State, with coastal south-west Western Australia experiencing the driest summer on record in 2023-2024 (BOM, 2025a). Summer rainfall was also below average in the west Kimberley, the Pilbara, and parts of the Southern Interior. Winter rainfall was mostly close to average in south-west Western Australia, whilst the north-west and central parts of the State recorded above average rainfall (BOM, 2025a; BOM, 2025b).

## *Western Grey Kangaroos*

Parts of the South-East PMZ have experienced average to above average rainfall and pasture growth over the last 12-24 months, however, the western half of the South-East PMZ near Esperance has continued to experience average to very much below average rainfall and extremely low pasture growth (Appendix 2). Population estimates for 2024 indicated that western grey kangaroos remained at a relatively low density of 0.24 individuals per km<sup>2</sup> across the South-East PMZ; falling below the no harvest threshold. Densities of western grey kangaroos were therefore calculated by KMA within the South-East PMZ (Table 1.4). The density of western grey kangaroos was 0.73 individuals per km<sup>2</sup> in the South Eastern Agricultural KMA and 0.45 individuals per km<sup>2</sup> in the Nullarbor KMA. Western grey kangaroo densities at the remaining KMA were below 0.3 individuals per km<sup>2</sup>. It is likely that higher quality habitat for western grey kangaroos exists in the coastal South Eastern Agricultural and Nullarbor KMA of the South-East PMZ (Figure 1, DBCA 2023).

In accordance with the Management Plan, Action 12 (DBCA 2023), in the South-East PMZ the commercial harvest rate is to be:

- reduced to 10% of the population estimate if western grey kangaroo population density estimates fall between 0.33 and 0.46 individuals per km<sup>2</sup>; or
- suspended if western grey kangaroo population density estimates fall below 0.33 individuals per km<sup>2</sup> within the South-East PMZ, or KMAs within the South-East PMZ.

Therefore, in the South-East PMZ in 2025, regular harvest of western grey kangaroos may occur only from the South Eastern Agricultural KMA and reduced harvest may occur from the Nullarbor KMA (Table 1.4). There is to be no harvest of western grey kangaroos in the remainder of the South-East PMZ in 2025. This is the fourth consecutive year in which harvest suspensions have been in place for the South-East PMZ, excluding the South Eastern Agricultural KMA where population densities have remained above the density thresholds. Regular harvest rates of 15% of the population estimate have been set for western grey kangaroos in the Central and South-West PMZs.

### *Red Kangaroos*

Population estimates for 2024 indicated that red kangaroo density (0.41 individuals per km<sup>2</sup>) in the Central PMZ has further reduced in response to ongoing below average rainfall and extremely low pasture growth over the last 24 months. In accordance with the Management Plan, Action 12 (DBCA, 2023), in the Central PMZ the commercial harvest rate is to be:

- reduced to 10% of the population estimate if red kangaroo density estimates fall between 0.40 and 0.57 individuals per km<sup>2</sup>.

Therefore, in the Central PMZ in 2025, a reduced harvest of red kangaroos may occur (Table 1.2).

In the Northern PMZ, 2024 population estimates indicate that the red kangaroo density (0.37 individuals per km<sup>2</sup>) has fallen into the reduced harvest rate category where the quota is to be set at 10% of the population estimate. The kangaroo density is very close to the no harvest threshold. Continued low rainfall and low pasture growth may mean this PMZ is closed in future years.

Regular harvest rates of 17% of the population estimate have been set for red kangaroos in the South-East PMZs.

Suspended harvests or reduced harvest rates will remain in place until surveys, or populations estimates corrected for trends in rainfall, indicate that kangaroo densities have increased above density thresholds.

## 1.1 State summary

Species	2025 Proposal			2024		
	2024 Population estimate	Harvest rate (%)	Quota	Quota	Harvest rate (%)	2023 Population estimate
Red kangaroo	527,850	13 <sup>a</sup>	<b>70,115</b>	89,330	15 <sup>a</sup>	609,155
Western grey kangaroo	776,240	14 <sup>a</sup>	<b>112,275</b>	161,100	15 <sup>a</sup>	1,086,235
<b>Totals</b>	<b>1,304,090</b>		<b>182,390</b>	250,430		<b>1,695,390</b>

<sup>a</sup> Rate calculated from total harvest quota

## 1.2 Regional quotas for red kangaroos in 2025

Population Monitoring Zone	2024 Population estimate ( $\hat{N}$ )	2025 Proposal	
		Harvest rate ( $H$ ) %	Quota ( $\hat{N} \times H$ )
Central	161,235	10	16,125
Northern	119,050	10	11,905
South-East	247,565	17	42,085
<b>Totals</b>	<b>527,850</b>		<b>70,115</b>


### 1.3 Regional quotas for western grey kangaroos in 2025

Population Monitoring Zone	2024 Population estimate ( $\hat{N}$ )	2025 Proposal	
		Harvest rate ( $H$ ) %	Quota ( $\hat{N} \times H$ )
Central	171,150	15	25,670
South-East	81,445	10-15 (NU and SEA Kangaroo Management Areas only)	8,060 (NU and SEA Kangaroo Management Areas only)
South-West	523,645	15	78,545
<b>Totals</b>	<b>776,240</b>		<b>112,275</b>


### 1.4 South-East Kangaroo Management Area quotas for western grey kangaroos in 2025

Kangaroo Management Area	2024 Density (kangaroos per km <sup>2</sup> )	2024 Population estimate ( $\hat{N}$ )	2025 Proposal	
			Harvest rate ( $H$ ) %	Quota ( $\hat{N} \times H$ )
South Eastern Agricultural (SEA)	0.73	29,155	15	4,375
Dundas (DU)	0.10	4,455	0	0
Nullarbor (NU)	0.45	36,840	10	3,685
Coolgardie (CO/CG)	0.03	1360	0	0
Leonora Eastern Goldfields (LEG)	0.07	7,585	0	0





*Figure 1: Kangaroo Population Monitoring Zones over the allowable harvest areas in Western Australia.*



*Figure 2: Kangaroo Management Areas in WA*

## 2 Aerial survey and population estimation methods

### 2.1 Fixed-wing aerial survey

The aerial survey program has been conducted across Western Australia since 1981 following pre-determined transect lines (Figure 1). The survey is conducted using standard 200 m fixed-width strip transect methods from a fixed-wing aircraft (Table 1; (Pople and Grigg, 1999). Correction factors are applied to account for unseen kangaroos (Table 2).

In 2024, the Central PMZ was scheduled to be surveyed in full as were the monitor blocks in the South-West PMZ (Table 1), however, due to safety concerns raised by pastoral lessees regarding potential interactions with mustering aircraft, all surveys were postponed for the year.

*Table 1 Aerial survey parameters*

<b>Altitude</b>	250 feet (76 m) above ground level
<b>Speed</b>	100 knots
<b>Strip width</b>	200 metres
<b>Survey unit</b>	5 kilometres long by 200 metres wide = 1 km <sup>2</sup>
<b>Survey line</b>	1 degree of longitude
<b>Survey intensity</b>	2 lines per one degree block (one degree latitude by one degree longitude)
<b>Species</b>	Red kangaroo ( <i>Osphranter rufus</i> ) and western grey kangaroo ( <i>Macropus fuliginosus</i> )
<b>Extent</b>	Pastoral rangelands and parts of the northern and south coast agricultural regions (Figure 1)
<b>Frequency</b>	Whole of commercial harvest zone was surveyed triennially from 1981 to 1993 (1981, 1984, 1987, 1990, 1993). Then: <ul style="list-style-type: none"> <li>Northern Zone in 1995, 1998, 2001, 2004, 2007, 2010, 2013, 2016, 2019 and 2022</li> <li>South-East Zone in 1996, 1999, 2002, 2005, 2008, 2011, 2014, 2017, 2021 and 2023</li> <li>Central Zone in 1997, 2000, 2003, 2006, 2009, 2012, 2015, 2018 and 2022</li> </ul>
<b>Monitor blocks</b>	Monitor blocks in the South-West Zone in 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2021, 2022 and 2023

<b>Methodology</b>	Standard 200 metre fixed-width strip surveys flown at 100 knots (185 km/hr), 76 metres above ground level (see Pople and Grigg 1999). Population estimates are made using standard correction factors (Table 2) with temperature correction for all species.
--------------------	--

*Table 2: Standard fixed-wing survey correction factors used for population estimates.*

Species	Correction Factor <sup>a</sup>			
	Open Vegetation	Light Vegetation	Medium Vegetation	Dense Vegetation
Red Kangaroo	2.29 <sup>b</sup>	2.36 <sup>b</sup>	2.43 <sup>b</sup>	2.57 <sup>b</sup>
Western Grey Kangaroo	4.8 <sup>c</sup>	4.8 <sup>c</sup>	4.8 <sup>c</sup>	4.8 <sup>c</sup>

<sup>a</sup> Used to correct raw counts to reliable population estimates. Accounts for the proportion of kangaroos missed by observers following standard counting methodologies.

<sup>b</sup> From Caughley, Sinclair and Scott-Kemmis (1976).

<sup>c</sup> From Pople and Grigg (1999).

## 2.2 Population estimates

The South-East and South-West PMZs were surveyed most recently in 2023 while the Northern and Central PMZs were surveyed most recently in 2022. No surveys were undertaken in 2024. Population estimates for all PMZs were determined in accordance with the management plan using the calculation for intervening survey years, i.e., the most recent population estimate adjusted for regional rainfall and commercial harvest offtake, according to the equation:

$$\hat{N}_{i+1} = (\hat{N}_i - H) \times r,$$

where:

$\hat{N}_i$  = the most recent population estimate;

$H$  = commercial harvest offtake between population estimates; and

$r$  = population growth rate for a regional rainfall category.

## 3 Criteria used in setting quotas

The proposed quota is set in accordance with Actions 9-12 of the Management Plan and takes into consideration information available on:

- Historical commercial harvest statistics (Appendix 1);
- Seasonal conditions (Appendix 2);
- Current populations trends (Appendix 3);

- The proportion of habitat and population not subject to harvesting;
- Current land use practice and trends in land use; and
- Significance of the non-commercial take relative to the population estimates, commercial quota and commercial harvest.

Quotas are based on a proportion threshold harvesting strategy as described in the Management Plan to ensure sustainable harvest (DBCA, 2023). Under this strategy, the quota does not exceed 17% of the estimated population of red kangaroos, and 15% of the estimated population of western grey kangaroos. Low populations are further protected by the implementation of density thresholds that trigger harvest reduction to 10% of the estimated population for both species, or harvest suspension. Thresholds have been determined based on standard deviations relative to the long-term average density and are specific to species and PMZ.

*Table 3: Density thresholds for Western Australia PMZs for red and western grey kangaroos.*

PMZ <sup>x</sup>	Density thresholds (kangaroos per km <sup>2</sup> )					
	Red kangaroo			Western grey kangaroo		
	17% HR (Threshold 1)	10% HR	No harvest (Threshold 2)	15% HR (Threshold 1)	10% HR	No harvest (Threshold 2)
Central	$D > 0.57^a$	$0.57^a \geq D > 0.40^b$	$D \leq 0.40^b$	$D > 0.18^a$	$0.18^a \geq D > 0.12^b$	$D \leq 0.12^b$
Northern	$D > 0.47^b$	$0.47^b \geq D > 0.36^c$	$D \leq 0.36^c$	Extralimital – no quota		
South-East	$D > 0.29^b$	$0.29^b \geq D > 0.23^c$	$D \leq 0.23^c$	$D > 0.46^b$	$0.46^b \geq D > 0.33^c$	$D \leq 0.33^c$
South-West	Vagrant – no quota			$D > 8.4^c$	$8.4^c \geq D > 6.5^d$	$D \leq 6.5^d$

<sup>x</sup> = The density thresholds for each PMZ will also apply to any Kangaroo Management Area within that PMZ.

HR = Harvest rate as a proportion of the population estimate.

D = Density estimate (kangaroos per km<sup>2</sup>) from aerial surveys. Threshold densities were calculated as 1.0, 1.5, 2 or 3 standard deviations from the mean of density estimates for full surveys of a PMZ between 1995 and 2022.

a = 1.0 standard deviation (sd); b = 1.5 sd; c = 2.0 sd; d = 3.0 sd

## 4 Harvest monitoring

Trends in harvest data including commercial take, sex ratio and average carcass weight by sex are analysed across the commercial harvest zone (Appendix 1).

## References

Bureau of Meteorology (2025a) Western Australia in summer 2023-24, Seasonal Climate Summary for Western Australia. Available at <http://www.bom.gov.au/climate/current/season/wa/archive/202402.summary.shtml> [Accessed 2 January 2025].

Bureau of Meteorology (2025b) Western Australia in winter 2024, Seasonal Climate Summary for Western Australia. Available at <http://www.bom.gov.au/climate/current/season/wa/archive/202408.summary.shtml> [Accessed 2 January 2025].

Caughley, G., Sinclair, R. and Scott-Kemmis, D. (1976) 'Experiments in aerial survey', *Journal of Wildlife Management*, 40(2), pp. 290–300.


Department of Biodiversity, Conservation and Attractions (2023) *Management plan for the commercial harvest of kangaroos in Western Australia 2024-2028*. Western Australia.

Pople, T. and Grigg, G. (1999) *Commercial harvesting of kangaroos in Australia*. Canberra: Environment Australia.


# Appendices

## Appendix 1 Harvest monitoring results for Western Australia


The figures and tables in Appendix 1 only include harvest data processed prior to 31 October 2024.




**Fig. A1.1.** Total commercial kangaroo harvest in Western Australia from 1983 to 2024. Commercial harvest of euros (*Osphranter robustus*) did not take place between 2003-2006 and from 2010-2024.



**Fig. A1.2.** Proportion of the commercial quota of red and western grey kangaroos harvested in Western Australia from 1972 to 2024.




**Fig. A1.3.** Sex ratio of the commercial kangaroo harvest in Western Australia from 1972 to 2024.




**Fig. A1.4.** Number and sex of red kangaroos harvested commercially in Western Australia from 1972 to 2024.






**Fig. A1.5.** Average carcass weights for red kangaroos harvested commercially in Western Australia from 1995 to 2024. Carcass dressing methods (and therefore carcass weights) are not standardised.




**Fig. A1.6.** Number and sex of western grey kangaroos harvested commercially in Western Australia from 1983 to 2024.




**Fig. A1.7.** Average carcass weights for western grey kangaroos harvested commercially in Western Australia from 1995 to 2024. Carcass dressing methods (and therefore carcass weights) are not standardised.


## Appendix 2 Rainfall and drought maps




**Fig. A2.1.** Rainfall deciles for Western Australia for the period 1 November 2023 to 31 October 2024 (last 12 months).




**Fig. A2.2.** Rainfall deciles for Western Australia for the period 1 November 2022 to 31 October 2024 (last 24 months).




**Fig. A2.3.** Drought map for Western Australia for the period 1 November 2023 to 31 October 2024 (last 12 months).




**Fig. A2.4.** Drought map for Western Australia for the period 1 November 2022 to 31 October 2024 (last 24 months).




**Fig. A2.5.** Rainfall totals for Western Australia for the period 1 November 2023 to 31 October 2024 (last 12 months).



**Fig. A2.6.** Rainfall totals for Western Australia for the period 1 November 2022 to 31 October 2024 (last 24 months).




**Fig. A2.7.** Summer rainfall deciles for Western Australia for the period 1 November 2023 to 30 April 2024.



**Fig. A2.8.** Winter rainfall deciles for Western Australia for the period 1 May 2024 to 31 October 2024.

**Pasture Growth Percentile**  
Relative to Historical Records from 1957 to 2024  
November 2023 to October 2024




Pasture growth includes grasses and crops

[www.LongPaddock.qld.gov.au](http://www.LongPaddock.qld.gov.au)

**Fig. A2.9.** Pasture growth in Western Australia for the period November 2023 to October 2024 (last 12 months).

**Pasture Growth Percentile**  
Relative to Historical Records from 1957 to 2024  
November 2022 to October 2024



Pasture growth includes grasses and crops


[www.LongPaddock.qld.gov.au](http://www.LongPaddock.qld.gov.au)

**Fig. A2.10.** Pasture growth in Western Australia for the period November 2022 to October 2024 (last 24 months).

## Appendix 3 Regional population estimates

Harvest data for 2024 presented in the figures and tables in Appendix 3 only includes data processed prior to 31 October 2024.

### Northern Population Management Zone



**Fig. A3.1.1.** Population estimates for red kangaroos in the Northern PMZ. Note, all estimates use standard habitat correction factors. Temperature corrections are applied to post-1993 data. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and scaled according to trends in rainfall.

**Table A3.1.1. Red kangaroo population estimates for the Northern Population Management Zone.**


Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2013	203,820±35,588	9,789	Average	1.1
2014	213,434	7,435	Above average	1.2
2015	247,200	6,755	Average	na
2016	502,800±37,100	2,495	Average	1.1
2017	550,340	1,561	Average	1.1
2018	603,660	3334	Very much below average	na
2019	313,850±27,200	4311	Below average	0.8
2020	247,630	1,562	Above average <sup>d</sup>	1.2
2021	295,280	0	Above average	1.2
2022 <sup>b</sup>	221,850±22,555	3761	Below average	0.8
2023	174,470	4400 <sup>c</sup>	Below average <sup>d</sup>	0.7
2024	119,050			

<sup>a</sup>  $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$  where:  $\hat{N}_t$  = the most recent population estimate;  $H$  = commercial harvest off-take between population estimates; and,  $r$  = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan.


<sup>b</sup> The most recent full survey of the Northern PMZ was flown in August 2022.

<sup>c</sup> The commercial harvest off-take in the Northern PMZ between 1 January 2024 and 31 October 2024.

<sup>d</sup> Rainfall in the Northern Zone for the preceding 12 months was considered to be below average for the zone overall (Figs A2.1 – A2.10).




**Fig. A3.1.2.** Proportion of the Northern Population Management Zone commercial quota harvested from 2008 to 2024.




**Fig. A3.1.3.** Sex ratio of the commercial red kangaroo harvest in the Northern Population Management Zone from 1972 to 2024.






**Fig. A3.1.4.** Average carcass weights of the commercial red kangaroo harvest in the Northern Population Management Zone from 1972 to 2024. Carcass dressing methods (and therefore carcass weights) are not standardised.

### Central Population Management Zone



**Fig. A3.2.1.** Population estimates for red kangaroos in the Central Population Management Zone of Western Australia. Note, all estimates use standard habitat and temperature correction factors. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and/or monitor block surveys, and are scaled according to trends in regional rainfall.



**Fig. A3.2.2.** Population estimates for western grey kangaroos in the Central Population Management Zone of Western Australia. Note, all estimates use standard habitat and temperature correction factors. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and/or monitor block surveys and are scaled according to trends in regional rainfall.

**Table A3.2.1. Red kangaroo population estimates for the years following a full aerial survey of the Central Population Management Zone.**

Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2012	141,765	7,333	Average	1.1
2013	147,875	10,704	Below average	0.8
2014	109,737	12,842	Above average	na
2015	485,000±29,000	7,886	Average	1.1
2016	524,800	3,399	Average	1.1
2017	573,540	2,599	Average	na
2018	857,350±64,300	3,335	Below average	0.8
2019	683,210	4,992	Below average	0.8
2020	542,575	2,621	Average	1.1
2021	593,950	1925	Above average	1.2
2022 <sup>b</sup>	341,870	3181	Very much below average	0.6
2023	203,210	1666 <sup>c</sup>	Below average <sup>d</sup>	0.8
2024	161,235			

<sup>a</sup>  $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$  where:  $\hat{N}_t$  = the most recent population estimate;  $H$  = commercial harvest off-take between population estimates; and,  $r$  = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan.

<sup>b</sup> The most recent full survey of the Central PMZ was flown in September 2022.

<sup>c</sup> The commercial harvest off-take in the Central PMZ between 1 January 2024 and 31 October 2024.

<sup>d</sup> Rainfall in the Central PMZ for the preceding 12 months was considered to be below average for the zone overall (Figs A2.1 – A2.10).

*Table A3.2.2. Western grey kangaroo population estimates for the Central Population Management Zone.*


Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2012	51,193	5,178	Average	1.1
2013	50,616	4,716	Below average	0.8
2014	36,720	5,504	Above average	na
2015	300,100±72,500	5,631	Average	1.1
2016	323,900	3,313	Average	1.1
2017	352,645	2,762	Average <sup>d</sup>	na
2018	637,660±157,800	2,679	Below average	0.8
2019	507,985	5,219	Below average <sup>d</sup>	0.8
2020	402,210	4,632 <sup>c</sup>	Average	1.1
2021	437,340	3,825	Above average	1.2
2022 <sup>b</sup>	369,330	4807	Very much below average	0.6
2023	218,710	4778 <sup>c</sup>	Below average <sup>d</sup>	0.8
2024	171,150			

<sup>a</sup>  $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$  where:  $\hat{N}_t$  = the most recent population estimate;  $H$  = commercial harvest off-take between population estimates; and,  $r$  = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan.

<sup>b</sup> The most recent full survey of the Central PMZ was flown in September 2022.

<sup>c</sup> The commercial harvest off-take in the Central PMZ between 1 January 2024 and 31 October 2024.


<sup>d</sup> Rainfall in the Central PMZ for the preceding 12 months was considered to be below average for the zone overall (Figs A2.1 – A2.10).



**Fig. A3.2.3.** Proportion of the Central Population Management Zone commercial quota harvested from 2008 to 2024.




**Fig. A3.2.4.** Sex ratio of the commercial red and western grey kangaroo harvest in the Central Population Management Zone from 2007 to 2024.




**Fig. A3.2.5.** Average carcass weights of the commercial red and western grey kangaroo harvest in the Central Zone from 2007 to 2024. Carcass dressing methods (and therefore carcass weights) are not standardised.

### South-East Population Monitoring Zone



**Fig. A3.3.1.** Population estimates for red kangaroos in the South-East Population Management Zone of Western Australia. Note, all estimates use standard habitat and temperature correction factors. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and/or monitor block surveys and are scaled according to trends in regional rainfall. \*\*The 2020 aerial survey was not conducted due to Covid-19 restrictions.



**Fig. A3.3.2.** Population estimates for western grey kangaroos in the South-East Population Management Zone of Western Australia. Note, all estimates use standard habitat and temperature correction factors. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and/or monitor block surveys and are scaled according to trends in regional rainfall. \*\*The 2020 aerial survey was not conducted due to COVID-19 restrictions.

**Table A3.3.1. Red kangaroo population estimates for the South-East Population Management Zone.**

Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2014	86,200±17,250	0	Average	1.1
2015	94,800	7,781	Above average	1.2
2016	104,400	5,580	Above average	na
2017	340,450±43,470	8,857	Average	1.1
2018	364,750	12,801	Below average	0.8
2019	281,560	14,452	Below average	0.8
2020	213,685	9,660	Average	1.1
2021	165,360±23,530	8,171	Average	1.1
2022	172,910	6,494	Below average	0.8
2023 <sup>b</sup>	231,475±37,515	6415 <sup>c</sup>	Average <sup>d</sup>	1.1
2024	247,565			
<sup>a</sup> $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$ where: $\hat{N}_t$ = the most recent population estimate; $H$ = commercial harvest off-take between population estimates; and, $r$ = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan. <sup>b</sup> The most recent full survey of the South-East PMZ was flown in August 2023. <sup>c</sup> The commercial harvest off-take in the South-East PMZ between 1 January 2024 and 31 October 2024. <sup>d</sup> Rainfall in the South-East Zone for the preceding 12 months was considered to be average for the zone overall (Figs A2.1 – A2.10).				



**Table A3.3.2. Western grey kangaroo population estimates for the South-East Population Management Zone.**


Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2014	179,400±51,751	14,743	Average	1.1
2015	181,100	9,679	Above average	1.2
2016	205,700	4,304	Above average	na
2017	747,700±93,400	5,117	Average	1.1
2018	816,840	5,548	Below average	0.8
2019	647,620	2,919	Below average	0.8
2020	515,760	3,820	Average	1.1
2021	115,895±20,760	506	Average	1.1
2022	126,930	1,771	Below average	0.8
2023 <sup>b</sup>	77,255±18250	3213 <sup>c</sup>	Average <sup>d</sup>	1.1
2024	81,445			

<sup>a</sup>  $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$  where:  $\hat{N}_t$  = the most recent population estimate;  $H$  = commercial harvest off-take between population estimates; and,  $r$  = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan.


<sup>b</sup> The most recent full survey of the South-East PMZ was flown in August 2023.

<sup>c</sup> The commercial harvest off-take in the South-East PMZ between 1 January 2024 and 31 October 2024.



<sup>d</sup> Rainfall in the South-East PMZ for the preceding 12 months was considered to be average for the zone overall (Figs A2.1 – A2.10).



**Fig. A3.3.3.** Proportion of the South-East Population Management Zone commercial quota harvested from 2008 to 2024. Note, no red kangaroos were harvested in 2015.




**Fig. A3.3.4.** Sex ratio of the commercial harvest of red and western grey kangaroos in the South-East Population Management Zone from 1972 to 2024. Note, no red kangaroos were harvested in 2015.



**Fig A3.3.5.** Average carcass weights of the commercial red and western grey kangaroo harvest in the South-East Population Management Zone. Carcass dressing methods (and therefore carcass weights) are not standardised.

### South-West Population Management Zone



**Fig. A3.4.1.** Aerial survey population estimates with standard errors for western grey kangaroos in South-West Population Management Zone monitor blocks from 2004-2024. Note, all estimates use standard habitat and temperature correction factors. No aerial surveys were undertaken in the South-West PMZ in 2011, 2019, 2020 and 2024. New transects were added in 2013, which increased the aggregate area of the monitor blocks. Consequently, adjustments have been made to population estimates in prior years. \*Estimates for years where the PMZ was not surveyed in full are based on estimates from surveys in previous years and/or monitor block surveys and are scaled according to trends in regional rainfall.

Note that population estimates for the South-West PMZ are a product of the mean kangaroo density in the monitor blocks and the aggregate area of the monitor blocks. No additional allowance is made for kangaroos occupying the unsurveyed portion of the South-West PMZ (i.e. for the purpose of the quota calculation, the density in the unsurveyed areas is treated as zero). However, western grey kangaroos are harvested in the unsurveyed parts of the South-West PMZ.

**Table A3.4.1. Western grey kangaroo population estimates for the South-West Population Management Zone.**


Year	Population estimate <sup>a</sup> ( $\hat{N}_t$ )	Commercial harvest off-take ( $H$ )	Zone Rainfall Category	Population growth rate ( $r$ )
2018	969,300±105,250	31,617	Below average	0.7
2019	656,380	43,645 <sup>c</sup>	Below average	0.8
2020	490,190	36,874	Above average <sup>d</sup>	1.2
2021	626,310±140,240	38,685	Average	1.1
2022	923,660±161,780	43,197	Below average	0.7
2023 <sup>b</sup>	790,270±147,518	42,211 <sup>c</sup>	Below average <sup>d</sup>	0.7
2024	523,643			

<sup>a</sup>  $\hat{N}_{t+1} = (\hat{N}_t - H) \times r$  where:  $\hat{N}_t$  = the most recent population estimate;  $H$  = commercial harvest off-take between population estimates; and,  $r$  = population growth rate for a regional rainfall category in accordance with Action 10 of the Management Plan.


<sup>b</sup> The most recent full survey of the South-West PMZ was flown in October 2023.

<sup>c</sup> The commercial harvest off-take in the South-West PMZ between 1 January 2024 and 31 October 2024.


<sup>d</sup> Rainfall in the South-West Zone for the preceding 12 months was considered to be below average for the zone overall (Figs A2.1 – A2.10).



**Fig. A3.4.2.** Proportion of the South-West Population Management Zone regional commercial quota harvested from 2008 to 2024.



**Fig. A3.4.3.** Sex ratio of the commercial harvest of western grey kangaroos in the South-West Population Management Zone from 1983 to 2024.



**Fig A3.4.4.** Average carcass weights of the commercial western grey kangaroo harvest in the South-West Population Management Zone. Carcass dressing methods (and therefore carcass weights) are not standardised.